
BOLLETTINO UNIONE MATEMATICA ITALIANA

BLAGOJ S. POPOV

On ultraspherical polynomials.

Bollettino dell'Unione Matematica Italiana, Serie 3, Vol. 14
(1959), n.1, p. 105–108.

Zanichelli

<http://www.bdim.eu/item?id=BUMI_1959_3_14_1_105_0>

L'utilizzo e la stampa di questo documento digitale è consentito liberamente per motivi di ricerca e studio. Non è consentito l'utilizzo dello stesso per motivi commerciali. Tutte le copie di questo documento devono riportare questo avvertimento.

*Articolo digitalizzato nel quadro del programma
bdim (Biblioteca Digitale Italiana di Matematica)
SIMAI & UMI
<http://www.bdim.eu/>*

On ultraspherical polynomials.

Nota di BLAGOJ S. POPOV (a Skopje, Yougoslavie)

Summary - *This paper will give several results involving the ultraspherical polynomials. The explicit expression for the product of these polynomials as a sum of such polynomials will be obtained. In particular Bailey's formula for the product of two associated Legendre functions is proved. Besides, some integrals involving ultraspherical polynomials have been evaluated.*

1. In hypergeometric notation, the ultraspherical polynomials may be defined by

$$(1) \quad P_n^{(\alpha, \alpha)}(x) = \frac{(1 + \alpha)_n}{n!} {}_2F_1 \left(-n, n + 2\alpha + 1, \alpha + 1, \frac{1-x}{2} \right),$$

with

$$(a)_n = \frac{\Gamma(a+n)}{\Gamma(a)} = a(a+1) \dots (a+n-1),$$

$$(a)_0 = 1.$$

Writing

$$\begin{aligned} (2) \quad {}_2F_1 \left(-m, m + 2\beta + 1, \beta + 1, \frac{1-x}{2} \right) {}_2F_1 \left(-n, n + 2\alpha + 1, \alpha + 1, \frac{1-x}{2} \right) \\ = \sum_{k=0}^{\infty} a_k {}_2F_1 \left(-m-n+2k, m+n+2\alpha-2k+1, \alpha+1, \frac{1-x}{2} \right), \end{aligned}$$

we evaluate the coefficients a_k by a direct method.

We find

$$\begin{aligned} a_k = \frac{(n+\alpha-k+1)_m (n+2\alpha+1)_{m-k} (m+2\beta+1)_m (-n)_k (-m)_k (\beta+1/2)_k}{(2n+2\alpha-2k+1)_{2m} (2m+2\beta-2k+1)_{2k} (\beta+1)_{m-k} 2^{-k} k!} \cdot \\ \cdot \frac{2m+2n+2\alpha-4k+1}{2m+2n+2\alpha-2k+1} {}_4F_3 \left[\begin{matrix} \beta-\alpha, k-m, 1/2-\alpha, -k, \\ n-k+1, k-m-n-2\alpha, 1/2+\beta \end{matrix} \middle| 1 \right], \end{aligned}$$

where

$${}_4F_3 \left[\begin{matrix} \alpha, \beta, \gamma, \delta \\ a, b, c \end{matrix} \middle| x \right] = \sum_{i=0}^{\infty} \frac{(\alpha)_i (\beta)_i (\gamma)_i (\delta)_i}{(a)_i (b)_i (c)_i} \frac{x^i}{i!}.$$

$$m \leq n.$$

From (1) and (2) we have the formula

$$\begin{aligned} (3) \quad & P_n^{(\alpha, \alpha)}(x) P_m^{(\beta, \beta)}(x) = \\ & = (m+2\beta+1)_m \sum_{k=0}^m \binom{m+n-2k}{n-k} \cdot \\ & \cdot \frac{(n+\alpha-k+1)_k (m+\beta-k+1)_k (m+n+\alpha-2k+1)_k}{(2m+2\beta-2k+1)_{2k} (2n+2\alpha-2k+1)_{2m} 2^{-2k} k!} \cdot \\ & \cdot (n+2\alpha+1)_{m-k} (\beta + 1/2)_k \frac{2m+2n+2\alpha-4k+1}{2m+2n+2\alpha-2k+1} \cdot \\ & \cdot {}_4F_3 \left[\begin{matrix} \beta-\alpha, k-m, 1/2-\alpha, -k \\ n-k+1, k-m, n-2\alpha, 1/2+\beta \end{matrix} \middle| 1 \right] P_{m+n-2k}^{(\alpha, \alpha)}(x), \end{aligned}$$

which give the composition of the ultraspherical polynomials.

When $\alpha = \beta$, (3) becomes

$$\begin{aligned} (4) \quad & P_n^{(\alpha, \alpha)}(x) P_m^{(\alpha, \alpha)}(x) = \\ & = (m+2\alpha+1)_m \sum_{k=0}^m \binom{m+n-2k}{n-k} \cdot \\ & \cdot \frac{(n+\alpha-k+1)_k (m+\alpha-k+1)_k (m+n+\alpha-2k+1)_k}{(2m+2\alpha-2k+1)_{2k} (2n+2\alpha-2k+1)_{2m} 2^{-2k} k!} \cdot \\ & \cdot (n+2\alpha+1)_{m-k} (\alpha + 1/2)_k \frac{2m+2n+2\alpha-4k+1}{2m+2n+2\alpha-2k+1} P_{m+n-2k}^{(\alpha, \alpha)}(x). \end{aligned}$$

If $\alpha = \beta = 0$, the formula (3) reduces to the well-known NEUMANN-ADAMS formula for LEGENDRE polynomials [1].

2. The connexion between the function $P_n^{(\alpha, \alpha)}(x)$ and the associated LEGENDRE function $P_n^r(x)$ is given by the equation

$$\begin{aligned} (5) \quad & P_{n-r}^{(r, r)}(x) = \frac{n! 2^r}{(n+r)!} (1-x^2)^{-r/2} P_n^r(x). \\ & (r \text{ is an integer}). \end{aligned}$$

Substituting $P_n^{(r,r)}(x)$ from (5) in (4) we get the remaining BAILEY's formula [2]

$$\begin{aligned} & (1-x^2)^{-r/2} P_n^{(r,r)}(x) P_m^{(r,r)}(x) = \\ & = 2^r \sum_{k=0}^{m-r} \frac{A_{m-k}^{-r} A_{k,-r} A_{n-k}^{-r}}{A_{m+n-r-k}^r} \frac{(m+n-2r-2k)!}{(m+n-2k)!} \cdot \\ & \cdot \frac{2m+2n-2r-4k+1}{2m+2n-2r-2k+1} P_{m+n-r-2k}^r(x), \end{aligned}$$

with

$$A_m^r = \frac{\left(\frac{1}{2}\right)_m}{(m+r)!}, \quad A_{k,r} = \frac{\left(\frac{1}{2}-r\right)_k}{k},$$

$$m \leq n.$$

Furthermore, by means (3) one can deduce the formula for the product of two associated LEGENDRE functions of different degree and different order.

Really, we find

$$\begin{aligned} & (1-x^2)^{-r/2} P_m^{(r,r)}(x) P_n^{(s,s)}(x) = \\ & = 2^r \sum_{k=0}^{m-r} \frac{A_{m-k}^{-r} A_{k,-r} A_{n-k}^{-s}}{A_{m+n-r-k}^s} \frac{(m+n-r-s-2k)!}{(m+n-r+s-2k)!} \cdot \\ & \cdot \frac{2m+2n-2r-4k+1}{2m+2n-2r-2k+1} {}_4F_3 \left[\begin{matrix} r-s, k+r-m, 1/2-s, -k, \\ n-k-s+1, k+r-m-n+s, 1/2+r, 1 \end{matrix} \right] \cdot \\ & P_{m+n-r-2k}^s(x), \\ & (m-r \leq n-s). \end{aligned}$$

3. Using (3) we can evaluate the integral

$$\int_{-1}^1 P_n^{(\alpha, \alpha)}(x) P_m^{(\beta, \beta)}(x) dx.$$

Since

$$P_n^{(\alpha, \alpha)}(1) = (-1)^n P_n^{(\alpha, \alpha)}(-1) = \frac{(1+\alpha)_n}{n!},$$

and

$$\int_{-1}^1 P_n^{(\alpha, \beta)}(x) dx = \frac{1 + (-1)^n}{n + 2\alpha} \frac{2(\alpha)_{n+1}}{(n+1)!},$$

we have

$$\begin{aligned} & \int_{-1}^1 P_n^{(\alpha, \beta)}(x) P_m^{(\beta, \beta)}(x) dx = \\ & = 2K \sum_{k=0}^m \frac{(n+\alpha-k+1)_m (m+\beta-k+1)_k (n+2\alpha+1)_{m-k} (\beta+1/2)_k}{(2n+2\alpha-2k+1)_{2m} (2m+2\beta-2k+1)_{2k} (n-k)! (m-k)! 2^{-2k} k!} \cdot \\ & \quad \cdot \frac{(m+n-2k+1)^{-1}}{(m+n+2\alpha-2k)} \cdot \\ & \quad \cdot \frac{2m+2n+2\alpha-4k+1}{2m+2n+2\alpha-2k+1} {}_4F_3 \left[\begin{matrix} \beta-\alpha, k-m, 1/2-\alpha, -k, \\ n-k+1, k-m-n-2\alpha, 1/2+\beta, 1 \end{matrix} \right]. \end{aligned}$$

$$K = [1 + (-1)^{m+n}](\alpha)_{n+1} (m+2\beta+1)_m.$$

When $m = n$, and $\beta = 0$, we obtain

$$\int_{-1}^1 P_n(x) P_n^{(\alpha, \alpha)}(x) dx = \frac{(n+2\alpha+1)_n}{2^{n-1} (2n+1)!!}.$$

If $\beta = 0$ and α is a positive integer, then

$$\int_{-1}^1 P_m(x) P_n^{(\alpha, \alpha)}(x) dx = \binom{k+\alpha-1}{k} \frac{2^\alpha (m+n+2\alpha-1)!!}{(n+\alpha)_\alpha (m+n+1)!!},$$

$$n - m = 2k.$$

REFERENCES

- [1]. WHITTAKER, E. T. and WATSON G. N., *A Course of modern Analysis*, fourth edition, Cambridge, 1952.
- [2]. BAILEY W. N., *On the Product of two associated Legendre Functions*, *The Quarterly Journal of Mathematics*, Oxford series, vol. II, N. 41, (1941), pp. 30-36.