
BOLLETTINO UNIONE MATEMATICA ITALIANA

RICHARD BELLMAN

Oscillatroy and unimodal properties of solutions of second order linear differential equations.

Bollettino dell'Unione Matematica Italiana, Serie 3, Vol. 19
(1964), n.3, p. 306–310.

Zanichelli

<http://www.bdim.eu/item?id=BUMI_1964_3_19_3_306_0>

L'utilizzo e la stampa di questo documento digitale è consentito liberamente per motivi di ricerca e studio. Non è consentito l'utilizzo dello stesso per motivi commerciali. Tutte le copie di questo documento devono riportare questo avvertimento.

*Articolo digitalizzato nel quadro del programma
bdim (Biblioteca Digitale Italiana di Matematica)
SIMAI & UMI*

<http://www.bdim.eu/>

Oscillatory and unimodal properties of solutions of second order linear differential equations (*)

Nota di RICHARD BELLMAN (Santa Monica, California, U.S.A.) (**)

Summary. - *In previous papers, we have shown how to derive fundamental properties of the solutions of*

$$u'' + p(x)u = 0$$

directly from the fact that it is the Euler equation associated with the functional

$$J(u) = \int_0^a [u'^2 - p(x)u^2] dx.$$

Here we wish to demonstrate a Sturm oscillation theorem and the unimodal property of the solution in analogous fashion, under the assumption that $p(x) > 0$.

1. Introduction.

In two previous papers, we have shown how to derive fundamental properties of the solutions of

$$(1.1) \quad u'' + p(x)u = 0$$

directly from the fact that it is the Euler equation associated

(*) This research is sponsored by the United States Air Force under Project RAND - Contract No. AF 49 (638) - 700 monitored by the Directorate of Development Planning, Deputy chief of Staff, Research and Development, Hq USAF. Views or conclusions contained in this Memorandum should not be interpreted as representing the official opinion or policy of the United States Air Force.

(**) Pervenuta alla Segreteria dell'U. M. I. il 16 giugno 1964.

with the functional

$$(1.2) \quad J(u) = \int_0^a [u'^2 - p(x)u^2] dx,$$

[1], [2]. Here we wish to demonstrate a STURM oscillation theorem and the unimodal property of the solution in analogous fashion, under the assumption that $p(x) > 0$.

2. The variational formulation.

Consider the problem of minimizing

$$(2.1) \quad K(u) = \frac{\int_0^a u'^2 dx}{\int_0^a p(x)u^2 dx}$$

over all functions $u(x)$ satisfying the conditions $u(0) = u(a) = 0$, and such that the numerator and denominator are nonzero. This leads in the usual way to the equation

$$(2.2) \quad u'' + \lambda p(x)u = 0, \quad u(0) = u(a) = 0,$$

with the minimum equal to the smallest characteristic value. If the equation

$$(2.3) \quad u'' + p(x)u = 0, \quad u(0) = u(a) = 0,$$

with $p(x) > 0$, has a solution which is positive for $0 < x < a$, we know that $\lambda = 1$ is the smallest characteristic value and that $u(x)$ is, up to a normalization factor, the corresponding characteristic function. This result can be established without the use of Sturmian oscillation theorems, if we employ the PERRON-FROBENIUS-JENTSCH techniques of positive operators.

3. The Sturmian property.

Suppose that we are given the two equations

$$(3.1) \quad \begin{aligned} u'' + p(x)u &= 0, & u(0) &= 0, \\ v'' + q(x)v &= 0, & v(0) &= 0, \end{aligned}$$

with $q(x) \geq p(x) > 0$. We wish to demonstrate that the first zero of $v(x)$ cannot occur after the first zero of u .

We assume that the situations were the reverse, as indicated below (see Fig. 1). Then, from what has preceded,

$$(3.2) \quad 1 = \min_v \frac{\int_0^b v'^2 dx}{\int_0^b q(x)v^2 dx} < \frac{\int_0^b w'^2 dx}{\int_0^b q(x)w^2 dx},$$




Fig. 1

where w is another function defined over $[0, b]$. Choose $w(x)$ in the following way:

$$(3.3) \quad \begin{aligned} w(x) &= u(x), & 0 \leq x \leq a, \\ &= 0, & a \leq x \leq b. \end{aligned}$$

Then (3.2) yields

$$(3.4) \quad 1 < \frac{\int_0^a u'^2 dx}{\int_0^a q(x)u^2 dx} \leq \frac{\int_0^a u'^2 dx}{\int_0^a p(x)u^2 dx} = 1,$$

a contradiction.

4. A unimodal property.

Let us now demonstrate a familiar property of the first characteristic function $u(x)$, its unimodal character. Suppose that $u(x)$, the solution of

$$(4.1) \quad u'' + p(x)u = 0, \quad u(0) = u(a) = 0,$$



Fig. 2

with the property that $u(x) > 0$, $0 < x < a$, had the form Fig. 2. Let us obtain from this function a function which yields a smaller

value of $K(u)$, as defined in (2.1), in the fashion of Fig. 3. In the interval $[a_2, a_1]$, $v(x) = u(a_1)$, otherwise $v(x) = u(x)$.




Fig. 3

We see that

$$(4.2) \quad \int_0^a v'^2 dx < \int_0^a u'^2 dx,$$

$$\int_0^a p(x)v^2 dx > \int_0^a p(x)u^2 dx.$$

Hence,

$$(4.3) \quad K(v) < K(u),$$

a contradiction to the minimizing property of u .

5. Discussion.

One value of the foregoing technique is that it can be applied to study non linear ordinary and partial differential equations as well. Furthermore, we can study the properties of the higher characteristic functions by using relative minima.

REFERENCES

- [1] R. BELLMAN, *On the Nonnegativity of Solutions of the Heat equation*, «Boll. d. Unione Matematica Italiana», Vol. 12, 1957, pp. 520-523.
- [2] — —, *On Variation-diminishing Properties of Green's Functions*, «Boll. d. Unione Matematica Italiana», Vol. 16, 1961, pp. 154-166.