
BOLLETTINO UNIONE MATEMATICA ITALIANA

DANIELE FAENZI

Cohomology of Tango bundle on \mathbb{P}^5

*Bollettino dell'Unione Matematica Italiana, Serie 8, Vol. 9-B (2006),
n.2, p. 319–326.*

Unione Matematica Italiana

[<http://www.bdim.eu/item?id=BUMI_2006_8_9B_2_319_0>](http://www.bdim.eu/item?id=BUMI_2006_8_9B_2_319_0)

L'utilizzo e la stampa di questo documento digitale è consentito liberamente per motivi di ricerca e studio. Non è consentito l'utilizzo dello stesso per motivi commerciali. Tutte le copie di questo documento devono riportare questo avvertimento.

*Articolo digitalizzato nel quadro del programma
bdim (Biblioteca Digitale Italiana di Matematica)
SIMAI & UMI*

<http://www.bdim.eu/>

Cohomology of Tango Bundle on \mathbb{P}^5 .

DANIELE FAENZI (*)

Sunto. – Il fibrato di Tango è definito come pull-back del fibrato di Cayley C su una quadrica liscia Q_5 in \mathbb{P}^6 attraverso una funzione f definita in caratteristica 2 che fattorizza il morfismo di Frobenius φ . La coomologia di T è calcolata in termini di $S \otimes C$, $\varphi^*(C)$, $\text{Sym}^2(C)$ e C , che si studiano con il teorema di Borel–Bott–Weil.

Summary. – The Tango bundle T is defined as the pull-back of the Cayley bundle over a smooth quadric Q_5 in \mathbb{P}^6 via a map f existing only in characteristic 2 and factorizing the Frobenius φ . The cohomology of T is computed in terms of $S \otimes C$, $\varphi^*(C)$, $\text{Sym}^2(C)$ and C , which we handle with Borel–Bott–Weil theorem.

1. – The bundles on \mathbb{P}^5 and on Q_5 .

The well-known Hartshorne conjecture states, in particular, that there are no indecomposable rank-2 vector bundles on \mathbb{P}^n , when n is greater than 5. However, one of the few rank-2 bundles on \mathbb{P}^5 up to twist and pull-back by finite morphisms is the Tango bundle T first given in [Tan76]. Later Horrocks in [Hor78] and Decker Manolache and Schreyer in [DMS92] discovered that it can be obtained starting from Horrocks rank-3 bundle. In [Fae03] we proved that T is the pull-back of the Cayley bundle over a smooth quadric Q_5 in \mathbb{P}^6 . Anyway it only exists in characteristic 2.

Here we compute the cohomology of T applying an analogue of Borel–Bott–Weil theorem in positive characteristic. In section (1) we introduce the involved bundles and state the theorem, while in section (2) we give proofs and some more remarks. We make use of Macaulay2 computer algebra package, see [GS].

Let k be an algebraically closed field, and let Q_5 be a smooth 5-dimensional quadric hypersurface over k . On the coordinate ring $R(\mathbb{P}^5)$ we use variables x_i 's

(*) The author was partially supported by Italian MIUR funds.


2000 Mathematics Subject Classification: Primary 14F05; Secondary 14J60.

while on $R(Q_5)$ we use z_j 's.

$$Q_5 = \{z_0^2 + z_1z_2 + z_3z_4 + z_5z_6 = 0\} \subset \mathbb{P}^6$$

Both on \mathbb{P}^5 and Q_5 , we denote Chern classes by integers, meaning integral multiples of the positive generators in each degree of the Chow ring. Further, we call *intermediate cohomology group* any sheaf cohomology group of degree other than 0 or 5, i.e. the dimension of the ambient variety.

On $Q_5 = G_2/P(a_1)$ we have the Cayley bundle C , associated to the standard representation of the semisimple part of the parabolic group $P(a_1)$, where a_1 is the shortest root in the Lie algebra of the exceptional Lie group G_2 , having the following Dynkin diagram.



The bundle C is obtained as the irreducible $P(a_1)$ -homogeneous module with maximal weight $\lambda_2 - 2\lambda_1$. The bundle $C(2)$ (weight λ_2) is globally generated and we have $h^0(C(2)) = 14$. By virtue of [Fae03, Theorem (2)] we can define the Tango bundle as follows.

DEFINITION. – *Let $\text{char}(k) = 2$. For any odd n there exists a non-constant morphism $f: \mathbb{P}^n \rightarrow Q_n$. For $n = 5$ the pull-back $T = f^*(C(1))$ is a stable rank-2 bundle with $c_1(T) = 2$, $c_2(T) = 4$ and Hilbert polynomial:*

$$\chi(T(t)) = \frac{1}{60}t^5 + \frac{1}{3}t^4 + \frac{25}{12}t^3 + \frac{11}{3}t^2 - \frac{51}{10}t - 14$$

Therefore we have, for every $0 \leq p \leq 5$ and $t \in \mathbb{Z}$:

$$T \simeq T^*(2) \quad \text{and} \quad h^p(T(t)) = h^{5-p}(T(-t-8))$$

THEOREM 1. – *Given the Tango bundle T defined above, we have:*

$$h^0(T(t)) = \chi(T(t)) \quad \text{for } t \geq 3$$

Furthermore, the only nonvanishing intermediate cohomology of T is the following:

t	-10	-9	-8	-7	-6	-5	-4	-3	-2	-1	0	1	2
$h^1(T(t))$									1	7	14	13	1
$h^2(T(t))$								1					
$h^3(T(t))$						1							
$h^4(T(t))$	1	13	14	7	1								

In particular, the bundle $T' = T(-4)$ is the only twist of T for which all cohomology groups vanish.

Recall the definition of the spinor 4-bundle S over Q_5 . Since $Q_5 = \text{Spin}(7)/\mathbf{P}(\beta_1)$, where β_1 is the first root of $\text{Spin}(7)$, we define S as the bundle associated to the spin representation of the semisimple part $ss(\mathbf{P}(\beta_1)) = \text{Spin}(5)$. The bundle $S^\vee = S(1)$ is globally generated. By Beilinson's theorem, the bundle $S(1)$, extended by zero to \mathbb{P}^6 , has the minimal graded free resolution:

$$0 \longrightarrow \mathcal{O}_{\mathbb{P}^6}(-1)^8 \xrightarrow{B} \mathcal{O}_{\mathbb{P}^6}^8 \longrightarrow S(1) \longrightarrow 0$$

where now B , by the observations in [Bea00], is an antisymmetric matrix whose determinant is the equation of the quadric, to the power 4. This is done by the matrix:

$$(1) \quad B = \begin{pmatrix} 0 & 0 & 0 & -z_3 & 0 & -z_1 & z_5 & -z_0 \\ 0 & 0 & z_3 & 0 & z_1 & 0 & -z_0 & -z_6 \\ 0 & -z_3 & 0 & 0 & -z_5 & z_0 & 0 & -z_2 \\ z_3 & 0 & 0 & 0 & z_0 & z_6 & z_2 & 0 \\ 0 & -z_1 & z_5 & -z_0 & 0 & 0 & 0 & z_4 \\ z_1 & 0 & -z_0 & -z_6 & 0 & 0 & -z_4 & 0 \\ -z_5 & z_0 & 0 & -z_2 & 0 & z_4 & 0 & 0 \\ z_0 & z_6 & z_2 & 0 & -z_4 & 0 & 0 & 0 \end{pmatrix}$$

The bundle C is related to S in the following way. One computes $c_4(S^\vee) = 0$. So, given a nontrivial global section of S^\vee , we have a rank-3 bundle G defined by:

$$(2) \quad 0 \longrightarrow \mathcal{O} \xrightarrow{a} S^\vee \longrightarrow G \longrightarrow 0$$

It turns out that $c_3(G^\vee(1)) = 0$. One can prove that $G^\vee(1)$ has a unique section b and that the quotient by such b is isomorphic to $C(1)$ i.e. C is the cohomology of the monad:

$$(3) \quad \mathcal{O}(-1) \xrightarrow{b(-1)} S \xrightarrow{a^t} \mathcal{O}$$

The bundle C has rank 2 and Chern classes $(-1, 1)$. The only non-vanishing intermediate cohomology groups are $H^1(C) = H^4(C(-4)) = k$. All this is done in [Ott90] and follows easily from [Jan87, Proposition 5.4] in any characteristic.

2. – Proof of the theorem and further remarks.

First of all we need some properties of the map $f: \mathbb{P}^n \longrightarrow Q_n$. Recall that it is defined for $\text{char}(k) = 2$ and for any odd n as:

$$f(x_0 : \dots : x_n) = (x_0x_1 + \dots + x_{n-1}x_n : x_0^2 : \dots : x_n^2).$$

The map f is a finite morphism of degree 2^{n-1} and factors the Frobenius morphism φ as in the diagram:

$$\begin{array}{ccc} & Q_n & \xrightarrow{\varphi} Q_n \\ f \nearrow & \downarrow \pi & \nwarrow f \\ \mathbb{P}^n & \xrightarrow{\varphi} & \mathbb{P}^n \end{array}$$

where π the projection from $(1 : 0 : \dots : 0)$.

OBSERVATION 2. – Given the map f defined above, we have:

$$(4) \quad \begin{aligned} f_*(\mathcal{O}_{\mathbb{P}^5}) &= \mathcal{O} \oplus \mathcal{O}(-1)^{14} \oplus \mathcal{O}(-2) \\ f_*(\mathcal{O}_{\mathbb{P}^5}(1)) &= S \oplus \mathcal{O}^6 \oplus \mathcal{O}(-1)^6 \end{aligned}$$

PROOF. – Let $\mathcal{F} = f_*(\mathcal{O}_{\mathbb{P}^5})$, $\mathcal{G} = f_*(\mathcal{O}_{\mathbb{P}^5}(1))$. The map f is a 16:1 cover, because the Frobenius is 32:1 and the projection π is 2:1. Then \mathcal{F} and \mathcal{G} are rank-16 vector bundles, whose cohomology one can read from the Leray degenerate spectral sequence. Indeed since $R^i(f_*) = 0$ (for $i > 0$) we have:

$$(5) \quad \begin{aligned} H^i(Q_5, \mathcal{F}(t)) &= H^i(\mathbb{P}^5, f^*(\mathcal{O}_{Q_5}(t))) = H^i(\mathbb{P}^5, \mathcal{O}_{\mathbb{P}^5}(2t)) \\ H^i(Q_5, \mathcal{G}(t)) &= H^i(\mathbb{P}^5, f^*(\mathcal{O}_{Q_5}(1)(t))) = H^i(\mathbb{P}^5, \mathcal{O}_{\mathbb{P}^5}(2t+1)) \end{aligned}$$

for $0 \leq i \leq 5$ and every t . This says that \mathcal{F} and \mathcal{G} have no intermediate cohomology, hence by [Kap86] or [BGS87] they must decompose as sum of twisted spinor bundles $S(a)$ and line bundles (although actually Kapranov's setting is over \mathbb{C}). For \mathcal{G} this implies, by a computation on the Euler characteristic, that the only choice is the one stated. On the other hand for \mathcal{F} we have a priori two possibilities: either the one stated above either $S \oplus S(-1) \oplus \mathcal{O} \oplus \mathcal{O}(-1)^6 \oplus \mathcal{O}(-2)$.

Now the formula (5) says that the polynomial ring $R(\mathbb{P}^5)$ decomposes as a module over $R(Q_5)$ (under the action given by f) as $R(\mathbb{P}^5)_{\text{even}} \oplus R(\mathbb{P}^5)_{\text{odd}}$ where:

$$R(\mathbb{P}^5)_{\text{even}} = \bigoplus_{t \in \mathbb{Z}} H^0(\mathbb{P}^5, \mathcal{O}(2t)) \quad R(\mathbb{P}^5)_{\text{odd}} = \bigoplus_{t \in \mathbb{Z}} H^0(\mathbb{P}^5, \mathcal{O}(2t+1))$$

For \mathcal{F} we have to compute explicitly a presentation of the $R(Q_5)$ -module $R(\mathbb{P}^5)_{\text{even}}$. We need e_0 to generate $k = R(\mathbb{P}^5)_0$ and e_{ij} to generate the monomial $x_i x_j$ ($i \neq j$) in $R(\mathbb{P}^5)_2$, thus obtaining a map:

$$\Phi: R(Q_5) \oplus R(Q_5)(-1)^{15} \longrightarrow R(\mathbb{P}^5)_0 \oplus R(\mathbb{P}^5)_2$$

But the coordinate e_{45} is redundant, since $z_0 \Phi(e_0) + \Phi(e_{01}) + \Phi(e_{23}) = \Phi(e_{45})$. Now for $R(\mathbb{P}^5)_4$: the terms containing x_i^2 already lie in the image (got by the action of z_{i-1}), and in fact we just have to fix $x_0 x_1 x_2 x_3$ because, e.g. $x_0 x_1 x_2 x_4 = z_3 \Phi(e_{34}) + z_5 \Phi(e_{25}) + z_0 \Phi(e_{24})$ and $x_0 x_1 x_4 x_5 = x_0 x_1 x_2 x_3 + z_1 z_2 \Phi(e_0) + z_0 \Phi(e_{01})$. Thus we get a generator in degree 2 (and no syzygy); moreover

$x_0x_1x_2x_3x_4x_5 = z_1z_2\Phi(e_{23}) + z_3z_4\Phi(e_{01}) + z_0(x_0x_1x_2x_3)$, so that $R(\mathbb{P}^5)_6$, and in fact all $R(\mathbb{P}^5)_{\text{even}}$, is also covered.

The presentation can be in fact computed also for $R(\mathbb{P}^5)_{\text{odd}}$. In this case one finds as syzygy of a map $R(Q_5)^6 \oplus R(Q_5)(-1)^{14} \rightarrow R(\mathbb{P}^5)_{\text{odd}}$ the matrix B described in (1) giving the spinor bundle, thus getting again (4). \square

REMARK 3. – Observation (2) can probably be extended to any odd dimension. Here we only mention that for $f: \mathbb{P}^3 \rightarrow Q_3$ we get $f_*(\mathcal{O}_{\mathbb{P}^3}) = S \oplus \mathcal{O}_{Q_3} \oplus \mathcal{O}_{Q_3}(-1)$ and $f_*(\mathcal{O}_{\mathbb{P}^3}(1)) = \mathcal{O}_{Q_3}^4$ which can be computed by the presentation or by Euler characteristic.

Finally, one may notice that $\pi_*(S) = \mathcal{O}_{\mathbb{P}^5}(-1)^8$ and clearly $\pi_*(\mathcal{O}_{\mathbb{P}^5}(t)) = \mathcal{O}(t) \oplus \mathcal{O}(t-1)$ so that the extension $0 \rightarrow S(-1) \rightarrow \mathcal{O}(-1)^8 \rightarrow S \rightarrow 0$ splits after π_* (actually this holds in any characteristic). This agrees with the formula:

$$\varphi_*(\mathcal{O}_{\mathbb{P}^5}) = \mathcal{O} \oplus \mathcal{O}(-1)^{15} \oplus \mathcal{O}(-2)^{15} \oplus \mathcal{O}(-3)$$

$$\varphi_*(\mathcal{O}_{\mathbb{P}^5}(1)) = \mathcal{O}^6 \oplus \mathcal{O}(-1)^{20} \oplus \mathcal{O}(-2)^6$$

Next we compute the cohomology of $\text{Sym}^2 C$, $C \otimes C$, $S \otimes C$. First notice that if V is the $P(a_1)$ -representation associated to the G_2 -homogeneous bundle C (of weight $\lambda_2 - 2\lambda_1$), $\text{Sym}^2 V$ is not completely reducible. This is due to the fact that $\text{SL}(2, k)$ is not linearly reductive in positive characteristic, check [Nag62].

In a similar way, the homogeneous bundle $\text{Sym}^2 C$ corresponds to a $P(a_1)$ -representation of weight $2\lambda_2 - 4\lambda_1$ which a rank-2 subrepresentation. In fact, setting $C^{[2]} = \varphi^*(C)$, we obtain a nonsplitting G_2 -equivariant exact sequence:

$$(6) \quad 0 \rightarrow C^{[2]} \rightarrow \text{Sym}^2 C \rightarrow \mathcal{O}(-1) \rightarrow 0$$

We also have:

$$(7) \quad \begin{aligned} \varphi_*(\mathcal{O}_{Q_5}) &= \mathcal{O} \oplus \mathcal{O}(-1)^{20} \oplus \mathcal{O}(-2)^7 \oplus S(-1) \\ \varphi_*(\mathcal{O}_{Q_5}(1)) &= \mathcal{O}^7 \oplus \mathcal{O}(-1)^{20} \oplus \mathcal{O}(-2) \oplus S \end{aligned}$$

Now again by Borel–Bott–Weil theorem ([Jan87, Proposition 5.4]) we know $h^i(\text{Sym}^2 C(3)) = 0$ for all i , $\text{Sym}^2 C(4)$ is globally generated and:

$$h^0(\text{Sym}(t)) = \chi(\text{Sym}^2 C(t)) = \frac{1}{20} t^5 + \frac{3}{8} t^4 - \frac{27}{8} t^2 - \frac{81}{20} t \quad \text{for } t \geq 4$$

$$h^1(\text{Sym}^2 C(2)) = 14 \quad h^1(\text{Sym}^2 C(1)) = 7 \quad h^1(\text{Sym}^2 C(-1)) = 1$$

where in the above twists these are the only non-vanishing H^i 's. By Serre duality we only miss $H^2(\text{Sym}^2 C)$. Now since $R^i(\varphi_*) = 0$, by the degenerate Leray spectral sequence, (6) and (7) we get:

$$\begin{aligned} H^2(\text{Sym}^2 C) &= H^2(C^{[2]}) \stackrel{\text{Leray}}{=} H^2(C \otimes S(-1)) \stackrel{\text{Leray}}{=} \\ &= H^2(C^{[2]}(-1)) = H^2(\text{Sym}^2 C(-1)) \stackrel{\text{Bott}}{=} k \end{aligned}$$

and by the same argument $h^1(\text{Sym}^2 C) = 1$, while the remaining h^i are zero. The same procedure yields the following values of $h^i(S \otimes C(t))$:

$$(8) \quad \begin{aligned} h^1(S \otimes C) &= 1 & h^1(S \otimes C(1)) &= 6 \\ h^2(S \otimes C(-1)) &= 1 & h^3(S \otimes C(-2)) &= 1 \\ h^4(S \otimes C(-4)) &= 6 & h^4(S \otimes C(-5)) &= 1 \end{aligned}$$

Now we can prove Theorem 1.

PROOF OF THEOREM 1. – Since $R^i(f_*) = 0$ for $i > 0$ we compute cohomology over Q_5 and get:

$$(9) \quad \begin{aligned} H^i(\mathbb{P}^5, T(2t)) &= H^i(Q_5, f_*(\mathcal{O}_{\mathbb{P}^5}) \otimes C(1+t)) \\ H^i(\mathbb{P}^5, T(2t+1)) &= H^i(Q_5, f_*(\mathcal{O}_{\mathbb{P}^5}(1)) \otimes C(1+t)) \end{aligned}$$

The even part gives $H^i(\mathbb{P}^5, T(2t)) = H^i(Q_5, C(1+t)) \oplus H^i(Q_5, C(t))^{14} \oplus H^i(Q_5, C(t-1))$, while the odd part gives $H^i(\mathbb{P}^5, T(2t+1)) = H^i(Q_5, C(1+t))^6 \oplus H^i(Q_5, C(t))^6 \oplus H^i(Q_5, S \otimes C(1+t))$ and these are known by the above formulas. \square

REMARK 4. – The values $h^2(\text{Sym}^2 C) = h^1(\text{Sym}^2 C) = 1$ exhibit non-standard cohomology for the representation $\text{Sym}^2 V$. Indeed $3\lambda_2 - 3\lambda_1$ is singular $((3\lambda_2 - 3\lambda_1, 3a_1 + a_2) = 0)$ so standard Borel–Bott–Weil theorem (i.e. in characteristic 0) would give $h^i(\text{Sym}^2 C) = 0$. Of course, we would have no such sequence as (6).

Still, by tensoring the monad defining C by $C(t)$ we get:

$$\begin{aligned} 0 \longrightarrow G^\vee \otimes C(t) &\longrightarrow S \otimes C(t) \longrightarrow C(t) \longrightarrow 0 \\ 0 \longrightarrow C(t-1) &\longrightarrow G^\vee \otimes C(t) \longrightarrow C \otimes C(t) \longrightarrow 0 \end{aligned}$$

whence we derive the values of $h^i(S \otimes C(t))$ from those of $\text{Sym}^2 C$, because, if $\text{char}(k) \neq 2$, $C \otimes C = \text{Sym}^2 C \oplus \mathcal{O}(-1)$. This way we would get the same values as in (8) if $\text{char}(k) = 0$. Finally, one can compute on Macaulay2 the values of the cohomology of $C^{[2]}$ and check the correctness of the above result.

Now we can write the Beilinson table of the normalized $T(-1)$:

$$\begin{pmatrix} 0 & 0 & 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 7 \\ 0 & 0 & 0 & 0 & 0 & 0 \end{pmatrix}$$

Then we write $T(-1)$ as cohomology of a monad:

$$\mathcal{O}(-1) \oplus \Omega^4(4) \longrightarrow \Omega^2(2) \oplus \Omega^1(1) \longrightarrow \mathcal{O}^7$$

As Wolfram Decker pointed out to us, another way to get Tango's bundle is by Horrocks bundle \mathcal{H} in characteristic 2. Concretely, the Horrocks bundle splits extension $0 \rightarrow T(-1) \rightarrow \mathcal{H} \rightarrow \mathcal{O} \rightarrow 0$. This allows to compute the cohomology of \mathcal{H} in terms of T .

Moreover in [DMS92] one finds an explicit description of the maps in the Beilinson monad. This provides also a check of computations. Denoting with e_0, \dots, e_5 the canonical basis of E (the exterior algebra over $W = H^0(\mathbb{P}^5, \mathcal{O}(1))$), and using the natural isomorphism $\text{Hom}(\Omega^i(i), \Omega^j(j)) = \wedge^{i-j} W = E_{j-i}$, T is the cohomology of the maps α and β :

$$\beta = \begin{pmatrix} e_0 & e_4 e_5 \\ e_1 & e_3 e_5 \\ e_2 & e_3 e_4 \\ e_3 & e_1 e_2 \\ e_4 & e_0 e_2 \\ e_5 & e_0 e_1 \\ 0 & e_0 e_3 + e_1 e_4 + e_2 e_5 \end{pmatrix}$$

$$\alpha = \begin{pmatrix} e_0 e_1 e_2 + e_3 e_4 e_5 & e_0 e_1 e_3 e_4 + e_0 e_2 e_3 e_5 + e_1 e_2 e_4 e_5 & 0 \\ e_0 e_3 + e_1 e_4 + e_2 e_5 & e_0 e_1 e_2 + e_3 e_4 e_5 & e_1 e_2 e_4 e_5 \end{pmatrix}$$

Finally, using the equations for T , **Macaulay2** provides the following resolution:

$$0 \rightarrow \bigoplus_{\mathcal{O}(-8)}^{\mathcal{O}^7(-7)} \rightarrow \mathcal{O}^{49}(-6) \rightarrow \mathcal{O}^{98}(-5) \rightarrow \mathcal{O}^{76}(-4) \rightarrow \bigoplus_{\mathcal{O}^{14}(-2)}^{\mathcal{O}^7(-3)} \rightarrow T \rightarrow 0$$

Applying cohomology algorithms in **Macaulay2** developed by Decker Eisenbud and Schreyer one may also obtain a full table of the cohomology, which we write in **Macaulay2** notation:

$$(10) \quad \begin{array}{r} \text{total: } 573 \ 260 \ 92 \ 27 \ 14 \ 7 \ 2 \ 2 \ 7 \ 14 \ 27 \ 92 \ 260 \ 573 \\ -6: \ 573 \ 260 \ 91 \ 14 \ . \ . \ . \ . \ . \ . \ . \ . \ . \ . \\ -5: \ . \ . \ 1 \ 13 \ 14 \ 7 \ 1 \ . \ . \ . \ . \ . \ . \\ -4: \ . \ . \ . \ . \ . \ . \ 1 \ . \ . \ . \ . \ . \ . \\ -3: \ . \ . \ . \ . \ . \ . \ . \ 1 \ . \ . \ . \ . \ . \\ -2: \ . \ . \ . \ . \ . \ . \ . \ 1 \ 7 \ 14 \ 13 \ 1 \ . \ . \\ -1: \ . \ . \ . \ . \ . \ . \ . \ . \ . \ 14 \ 91 \ 260 \ 573 \end{array}$$

Reading the table along one antidiagonal gives the list of cohomology groups of a single twist. Here the list for T starts from the up-right corner, while

starting from a shift to the left means reading the list for a (-1) -twist. Table (10) agrees with Theorem 1.

REFERENCES

- [Bea00] ARNAUD BEAUVILLE, *Determinantal hypersurfaces*, Michigan Math. J. **48** (2000), 39-64.
- [BGS87] RAGNAR-OLAF BUCHWEITZ - GERT-MARTIN GREUEL - FRANK-OLAF SCHREYER, *Cohen-Macaulay modules on hypersurface singularities. II*, Invent. Math. **88** (1987), no. 1, 165-182.
- [DMS92] WOLFRAM DECKER - NICOLAE MANOLACHE - FRANK-OLAF SCHREYER, *Geometry of the Horrocks bundle on \mathbf{P}^5* , Complex projective geometry (Trieste, 1989/ Bergen, 1989), London Math. Soc. Lecture Note Ser., vol. 179, Cambridge Univ. Press, Cambridge, 1992, pp. 128-148.
- [Fae03] DANIELE FAENZI, *A Geometric Construction of Tango Bundle on \mathbf{P}^5* , Kodai Mathematical Journal **27** (2004), no. 1, 1-6.
- [GS] DANIEL R. GRAYSON - MICHAEL E. STILLMAN, *Macaulay 2, a software system for research in algebraic geometry*, Available at <http://www.math.uiuc.edu/Macaulay2/>.
- [Hor78] GEOFFREY HORROCKS, *Examples of rank three vector bundles on five-dimensional projective space*, J. London Math. Soc. (2) **18** (1978), no. 1, 15-27.
- [Jan87] JENS CARSTEN JANTZEN, *Representations of algebraic groups*, Pure and Applied Mathematics, vol. 131, Academic Press Inc., Boston, MA, 1987.
- [Kap86] MIKHAIL M. KAPRANOV, *Derived category of coherent bundles on a quadric*, Funktsional. Anal. i Prilozhen. **20** (1986), no. 2, 67; English translation in Funct. Anal. Appl. **20** (1986), 141-142.
- [Nag62] MASAYOSHI NAGATA, *Complete reducibility of rational representations of a matrix group*, J. Math. Kyoto Univ. **1** (1961/1962), 87-99.
- [Ott90] GIORGIO OTTAVIANI, *On Cayley bundles on the five-dimensional quadric*, Boll. Un. Mat. Ital. A (7) **4** (1990), no. 1, 87-100.
- [Tan76] HIROSHI TANGO, *On morphisms from projective space P^n to the Grassmann variety $Gr(n, d)$* , J. Math. Kyoto Univ. **16** (1976), no. 1, 201-207.

Dipartimento di Matematica U. Dini
 Università di Firenze, Viale Morgagni 67/a
 I-50134, Florence, Italy
 E-mail: faenzi@math.unifi.it
 home page: <http://www.math.unifi.it/faenzi/>