

---

ATTI ACCADEMIA NAZIONALE DEI LINCEI  
CLASSE SCIENZE FISICHE MATEMATICHE NATURALI  
**RENDICONTI**

---

S. S. SINGH

**Conharmonic transformations of Einstein-Kähler  
spaces with Bochner curvature tensor**

*Atti della Accademia Nazionale dei Lincei. Classe di Scienze Fisiche,  
Matematiche e Naturali. Rendiconti, Serie 8, Vol. **57** (1974), n.6, p. 559–564.  
Accademia Nazionale dei Lincei*

<[http://www.bdim.eu/item?id=RLINA\\_1974\\_8\\_57\\_6\\_559\\_0](http://www.bdim.eu/item?id=RLINA_1974_8_57_6_559_0)>

L'utilizzo e la stampa di questo documento digitale è consentito liberamente per motivi di ricerca e studio. Non è consentito l'utilizzo dello stesso per motivi commerciali. Tutte le copie di questo documento devono riportare questo avvertimento.

Atti della Accademia Nazionale dei Lincei. Classe di Scienze Fisiche, Matematiche e Naturali. Rendiconti, Accademia Nazionale dei Lincei, 1974.

**Geometria differenziale.** — *Conharmonic transformations of Einstein-Kähler spaces with Bochner curvature tensor.* Nota di S. S. SINGH, presentata (\*) dal Socio E. BOMPIANI.

**RIASSUNTO.** — L'Autore, che già si era occupato di spazi Einstein-Kähleriani di tipo speciale, si occupa ora di detti spazi nell'ipotesi che il tensore di curvatura di Bochner soddisfi a speciali relazioni.

### I. INTRODUCTION

An  $n (= 2m)$  dimensional Kähler space  $K''$  is a Riemannian space if it admits a structure tensor  $\varphi_\mu^\lambda$  satisfying

$$(1.1) \quad \varphi_\mu^\alpha \varphi_\alpha^\lambda = -\delta_\mu^\lambda$$

$$(1.2) \quad \varphi_{\lambda\mu} = -\varphi_{\mu\lambda}, \quad (\varphi_{\lambda\mu} = \varphi_\lambda^\alpha g_{\alpha\mu}),$$

and

$$(1.3) \quad \varphi_{\lambda}^k,_\mu = 0,$$

where the comma followed by an index denotes the operator of covariant differentiation with respect to the metric tensor  $g_{\lambda\mu}$  of the Riemannian space.

It is well known that in an Einstein-Kähler space the Bochner curvature tensor reduces to  $U_{\lambda\mu\nu}^k$  given by

$$(1.4) \quad U_{\lambda\mu\nu}^k = R_{\lambda\mu\nu}^k + \frac{R}{n(n+2)} [g_{\lambda\nu} \delta_\mu^k - g_{\mu\nu} \delta_\lambda^k + \varphi_{\lambda\nu} \varphi_\mu^k - \varphi_{\mu\nu} \varphi_\lambda^k + 2 \varphi_{\lambda\mu} \varphi_\nu^k].$$

A Kähler space is called a space of constant holomorphic sectional curvature if the tensor  $U_{\lambda\mu\nu}^k$  given by (1.4) vanishes identically.

We shall consider Einstein-Kähler spaces which we shall denote by  $E''$ , satisfying

$$(1.5) \quad U_{\lambda\mu\nu}^k,_\epsilon = 0$$

and

$$(1.6) \quad U_{\lambda\mu\nu}^k,_\epsilon = K_\epsilon U_{\lambda\mu\nu}^k,$$

where  $K_\epsilon$  is a non-zero vector. An Einstein-Kähler space satisfying (1.5) is called a Kähler space with parallel Bochner curvature tensor and that

(\*) Nella seduta del 14 dicembre 1974.

(1) Numbers in square brackets refer to the references given at the end.

(2) As for the notations we follow K.B. Lal and S. S. Singh [3] and S. S. Singh [4].

satisfying (1.6) is called a Kähler space with recurrent Bochner curvature tensor. For brevity, such Einstein-Kähler spaces shall be denoted by  $Bs(E'')$  and  $Br(E'')$  respectively.

## 2. CONHARMONIC TRANSFORMATION OF EINSTEIN-KÄHLER SPACES

Let  $E''$  and  $\bar{E}''$  be two Einstein Kähler spaces. If the metric tensor  $\bar{g}_{\lambda\mu}$  of  $\bar{E}''$  is given by

$$(2.1) \quad \bar{g}_{\lambda\mu} = e^{2\sigma} g_{\lambda\mu}$$

where  $g_{\lambda\mu}$  is the metric tensor of  $E''$ , then  $\bar{E}''$  is said to be a conformal transformation of  $E''$ .

By a conformal transformation (2.1) the tensor  $\varphi_{\lambda}^{\mu}$  remains unchanged, but the tensor  $\varphi_{\lambda\mu}$  is transformed into  $\bar{\varphi}_{\lambda\mu}$  by

$$(2.2) \quad \bar{\varphi}_{\lambda\mu} = e^{2\sigma} \varphi_{\lambda\mu}.$$

Also, we have

$$(2.3) \quad \bar{R}_{\lambda\mu\nu k} = e^{2\sigma} [R_{\lambda\mu\nu k} + g_{\mu\nu} \sigma_{\lambda k} + g_{\lambda k} \sigma_{\mu\nu} - g_{\nu k} \sigma_{\mu\nu} - g_{\mu k} \sigma_{\lambda\nu}]$$

and

$$(2.4) \quad \bar{R}_{\lambda\mu\nu}{}^k = R_{\lambda\mu\nu}{}^k + g_{\nu\mu} \sigma_{\lambda}{}^k - g_{\nu\lambda} \sigma_{\mu}{}^k + \delta_{\lambda}{}^k \sigma_{\nu\mu} - \delta_{\mu}{}^k \sigma_{\nu\lambda}.$$

A conformal transformation with  $\sigma$  satisfying

$$(2.5) \quad \sigma_{,\alpha} + \frac{1}{2}(n-2)\sigma_{\alpha}\sigma^{\alpha} = 0$$

where  $\sigma_{\alpha} = \sigma_{,\alpha}$  and  $\sigma^{\alpha} = g^{\alpha\beta} \sigma_{\beta}$ , has been called a conharmonic transformation and studied by Y. Ishii [1, 2]. A conharmonic transformation satisfies the condition

$$(2.6) \quad (3) \quad \bar{g}_{\lambda\mu} \bar{R} = g_{\lambda\mu} R$$

which, in view of (1.2) gives

$$(2.7) \quad \bar{\varphi}_{\lambda\mu} \bar{R} = \varphi_{\lambda\mu} R.$$

When an Einstein-Kähler space is transformed into an Einstein-Kähler space by a conharmonic transformation  $\sigma$  satisfies the condition Y. Ishii [1]

$$(2.8) \quad \sigma_{\mu\nu} = 0.$$

Using (2.8) in (2.4) we have

$$(2.9) \quad \bar{R}_{\lambda\mu\nu}{}^k = R_{\lambda\mu\nu}{}^k.$$

In view of (2.6), (2.7) and (2.9) we obtain

$$(2.10) \quad \bar{U}_{\lambda\mu\nu}{}^k = U_{\lambda\mu\nu}{}^k.$$

(3) Y. Ishii [1].

3. CONHARMONIC TRANSFORMATION OF  $B\gamma(E'')$  SPACES

Differentiating (2.10) covariantly and using the relation

$$\left\{ \begin{array}{l} \bar{\alpha} \\ \lambda\mu \end{array} \right\} = \left\{ \begin{array}{l} \alpha \\ \lambda\mu \end{array} \right\} + \delta_\lambda^\alpha \sigma_\mu + \delta_\mu^\alpha \sigma_\lambda - \sigma^\alpha g_{\lambda\mu},$$

we get

$$(3.1) \quad \begin{aligned} \bar{U}_{\lambda\mu\nu}^k;_\varepsilon &= U_{\lambda\mu\nu}^k,_\varepsilon - 2\sigma_\varepsilon U_{\lambda\mu\nu}^k - [U_{\lambda\mu\nu\varepsilon} \sigma^k + U_{\varepsilon\mu\nu}^k \sigma_\lambda + \\ &+ U_{\lambda\nu\varepsilon}^k \sigma_\mu + U_{\lambda\mu\varepsilon}^k \sigma_\nu] + \sigma^\alpha [U_{\lambda\mu\nu\alpha} \delta_\varepsilon^k + U_{\alpha\mu\nu}^k g_{\lambda\varepsilon} + U_{\lambda\alpha\nu}^k g_{\mu\varepsilon} + U_{\lambda\mu\alpha}^k g_{\nu\varepsilon}] \end{aligned}$$

where the semi colon denotes the covariant differentiation with respect to  $\bar{g}_{\lambda\mu}$ .

Now we assume that both  $E''$  and  $\bar{E}''$  are  $B\gamma(E'')$  spaces, so that we have

$$(3.2) \quad U_{\lambda\mu\nu}^k,_\varepsilon = K_\varepsilon U_{\lambda\mu\nu}^k$$

$$(3.3) \quad \bar{U}_{\lambda\mu\nu}^k;_\varepsilon = \bar{K}_\varepsilon \bar{U}_{\lambda\mu\nu}^k$$

for non-zero vectors  $K_\varepsilon$  and  $\bar{K}_\varepsilon$ .

Substituting (3.2) and (3.3) in (3.1) and using (2.10) we obtain

$$(3.4) \quad \begin{aligned} (\bar{K}_\varepsilon - K_\varepsilon) U_{\lambda\mu\nu}^k &= -2\sigma_\varepsilon U_{\lambda\mu\nu}^k - [U_{\lambda\mu\nu\varepsilon} \sigma^k + U_{\varepsilon\mu\nu}^k \sigma_\lambda + \\ &+ U_{\lambda\nu\varepsilon}^k \sigma_\mu + U_{\lambda\mu\varepsilon}^k \sigma_\nu] + \sigma^\alpha [U_{\lambda\mu\nu\alpha} \delta_\varepsilon^k + U_{\alpha\mu\nu}^k g_{\lambda\varepsilon} + U_{\lambda\alpha\nu}^k g_{\mu\varepsilon} + U_{\lambda\mu\alpha}^k g_{\nu\varepsilon}]. \end{aligned}$$

Contracting  $\varepsilon$  and  $k$  in (3.4) and using the relations

$$(3.5) \quad U_{\alpha\lambda\mu}^\alpha = U_{\lambda\alpha\mu}^\alpha = U_{\lambda\mu\alpha}^\alpha = 0$$

and

$$(3.6) \quad U_{\lambda\mu\nu}^k + U_{\mu\nu\lambda}^k + U_{\nu\lambda\mu}^k = 0$$

we obtain

$$(3.7) \quad (\bar{K}_\alpha - K_\alpha) U_{\lambda\mu\nu}^\alpha = (n-3)\sigma_\alpha U_{\lambda\mu\nu}^\alpha.$$

Transvecting (3.7) with  $\sigma^\nu$  we get

$$(3.8) \quad (\bar{K}_\alpha - K_\alpha) U_{\lambda\mu\beta}^\alpha \sigma^\beta = 0.$$

On the other hand transvection of (3.4) with  $\sigma^\varepsilon$  gives

$$(3.9) \quad (\bar{K}_\alpha - K_\alpha + 2\sigma_\alpha) U_{\lambda\mu\nu}^k \sigma^\alpha = 0.$$

Hence, we find either

$$(3.10) \quad U_{\lambda\mu\nu}^k = 0$$

or

$$(3.11) \quad (\bar{K}_\alpha - K_\alpha) \sigma^\alpha = -2\sigma_\alpha \sigma^\alpha.$$

We consider the case in which (3.11) holds. Transvecting (3.4) with  $(\bar{K}_k - K_k) \sigma^v$  we have

$$\begin{aligned} (\bar{K}_\varepsilon - K_\varepsilon)(\bar{K}_k - K_k) \sigma^v U_{\lambda\mu\nu}{}^k &= -2(\bar{K}_k - K_k) U_{\lambda\mu\nu}{}^k \sigma^v \sigma_\varepsilon - \\ &- \{(\bar{K}_k - K_k) U_{\lambda\mu\nu\varepsilon} \sigma^k \sigma^v + (\bar{K}_k - K_k) U_{\varepsilon\mu\nu}{}^k \sigma^v \sigma_\lambda + \\ &+ (\bar{K}_k - K_k) \sigma^v \sigma_\mu U_{\lambda\nu\varepsilon}{}^k + (\bar{K}_k - K_k) U_{\lambda\mu\varepsilon}{}^k \sigma_\nu \sigma^v\} + \\ &+ \sigma^\alpha \{(\bar{K}_\varepsilon - K_\varepsilon) U_{\lambda\mu\nu\alpha} \sigma^v + (\bar{K}_k - K_k) U_{\alpha\mu\nu}{}^k g_{\lambda\varepsilon} \sigma^v + \\ &+ (\bar{K}_k - K_k) U_{\lambda\nu\alpha}{}^k g_{\mu\varepsilon} \sigma^v + (\bar{K}_k - K_k) U_{\lambda\mu\alpha}{}^k g_{\nu\varepsilon} \sigma^v\} \end{aligned}$$

which on using equations (3.11), (3.7) and (3.8) reduces to (3.12)

$$(3.12) \quad (n - 1) \sigma_\alpha \sigma^\alpha U_{\lambda\mu\nu}{}^\beta \sigma_\beta = 0.$$

Hence, we find either

$$(3.13) \quad \sigma_\alpha \sigma^\alpha = 0,$$

which implies that  $\sigma$  is constant

or

$$(3.14) \quad \sigma_\beta U_{\lambda\mu\nu}{}^\beta = 0.$$

When (3.14) holds, (3.4) becomes

$$(\bar{K}_\varepsilon - K_\varepsilon) U_{\lambda\mu\nu}{}^k = -2 \sigma_\varepsilon U_{\lambda\mu\nu}{}^k - [U_{\lambda\mu\nu\varepsilon} \sigma^k + U_{\varepsilon\mu\nu}{}^k \sigma_\lambda + U_{\lambda\nu\varepsilon}{}^k \sigma_\mu + U_{\lambda\mu\varepsilon}{}^k \sigma_\nu].$$

Transvecting this equation with  $\sigma_k$  and using (3.14) we get

$$\sigma_k \sigma^k U_{\lambda\mu\nu\varepsilon} = 0.$$

Hence, we find either (3.10) or (3.13). In the case of (3.13) the equation (3.4) becomes

$$(\bar{K}_\varepsilon - K_\varepsilon) U_{\lambda\mu\nu}{}^k = 0,$$

from which follows either (3.10) or  $\bar{K}_\varepsilon = K_\varepsilon$ .

Thus we have

**THEOREM 3.1.** *If a  $Br(E'')$  is transformed into another  $Br(E'')$  by a conharmonic transformation, then the following cases occur:*

- 1) *the space is that of constant holomorphic sectional curvature.*
- 2)  *$\sigma$  is constant and the recurrence vectors coincide.*

Since an Einstein-Kähler recurrent space is a  $Br(E'')$ , from the above theorem we have the following

**COROLLARY.** *If an Einstein-Kähler space is transformed into another Einstein-Kähler space by a conharmonic transformation as follows:*

*a  $Br(E'')$ -space  $\rightarrow$  a recurrent  $E''$ -space  
or*

*a recurrent  $E''$ -space  $\rightarrow$  a  $Br(E'')$ -space or a recurrent  $E''$ -space  
then the space is that of constant holomorphic sectional curvature or  $\sigma$  is constant and the recurrence vectors coincide.*

Now, if  $\sigma$  is constant and the space is not of constant holomorphic sectional curvature, then (3.1) can be written as

$$\bar{U}_{\lambda\mu\nu}{}^k; \varepsilon = U_{\lambda\mu\nu}{}^k, \varepsilon.$$

Consequently, a  $Br(E'')$  may be transformed into a  $Br(E'')$  by a conharmonic transformation.

Therefore, in view of Theorem 3.1, we have

**THEOREM 3.2.** *The necessary and sufficient condition for a  $Br(E'')$  which is not of constant holomorphic sectional curvature to be transformed into another  $Br(E'')$  is that  $\sigma$  is constant.*

#### 4. CONHARMONIC TRANSFORMATION OF $Bs(E'')$ SPACES

If  $E''$  and  $\bar{E}''$  are both  $Bs(E'')$ , then regarding  $K_\varepsilon$  and  $\bar{K}_\varepsilon$  both zero and proceeding exactly in the same way as in the proof of Theorem 3.1, we find either  $U_{\lambda\mu\nu}{}^k = 0$  or  $\sigma$  is constant. Hence, we have

**THEOREM 4.1.** *If a  $Bs(E'')$  is transformed into another  $Bs(E'')$  by a conharmonic transformation, then the space is that of constant holomorphic sectional curvature or  $\sigma$  is constant.*

**COROLLARY.** *If an Einstein-Kähler space is transformed into another Einstein-Kähler space by a conharmonic transformation as follows:*

a  $Bs(E'')$ -space  $\rightarrow$  a symmetric  $E''$ -space

or

a symmetric  $E''$ -space  $\rightarrow$  a  $Bs(E'')$ -space or a symmetric  $E''$ -space

then the space is that of constant holomorphic sectional curvature or  $\sigma$  is constant.

Also we have

**THEOREM 4.2.** *A necessary and sufficient condition for a  $Bs(E'')$  which is not of constant holomorphic sectional curvature to be transformed into another  $Bs(E'')$  by a conharmonic transformation is that  $\sigma$  is constant.*

Next, we assume that an  $E''$  is a  $Br(E'')$  and  $\bar{E}''$  is a  $Bs(E'')$ . Then regarding  $\bar{K}_\varepsilon$  as zero and proceeding exactly the same way as in the proof of Theorem 3.1 we find either  $U_{\lambda\mu\nu}{}^k = 0$  or  $\sigma$  is constant and  $K_\varepsilon = \bar{K}_\varepsilon = 0$ . However, since  $K_\varepsilon$  is a non-zero vector, the space must be that of constant holomorphic sectional curvature. Hence, we have

**THEOREM 4.3.** *If a  $Br(E'')$  is transformed into a  $Bs(E'')$  or a  $Bs(E'')$  is transformed into a  $Br(E'')$  by a conharmonic transformation, then the space is that of constant holomorphic sectional curvature.*

Since a Kähler space which is symmetric in the sense of Cartan is a  $B_s(E^n)$ , from this theorem we have the following.

COROLLARY. *If an  $E^n$  is transformed into another  $E^n$  by a conharmonic transformation as follows:*

*a  $Br(E^n)$ -space  $\rightarrow$  a symmetric  $E^n$ -space*

*a  $B_s(E^n)$ -space  $\rightarrow$  a recurrent  $E^n$ -space*

*a recurrent  $E^n$ -space  $\rightarrow$  a  $B_s(E^n)$ -space or a symmetric  $E^n$ -space*

*a symmetric  $E^n$ -space  $\rightarrow$  a  $Br(E^n)$ -space or a recurrent  $E^n$ -space*

*then, the space must be that of constant holomorphic sectional curvature.*

The Author is thankful to prof. K. B. Lal for his helpful guidance.

#### REFERENCES

- [1] Y. ISHII (1957) – *On conharmonic transformations*, «Tensor, N. S.», 7 (2).
- [2] Y. ISHII, (1965) – *On conharmonic transformations in subspaces*, «Memoirs of the Konan Univ. Sci. ser.», 9 (37).
- [3] K. B. LAL and S. S. SINGH (1971) – *On Kählerian spaces with recurrent Bochner curvature*, «Rend. Accad. Naz. dei Lincei» (8), LI, 213–220.
- [4] S. S. SINGH (1973) – *Some theorems on Kählerian spaces with parallel Bochner curvature tensor*, «Rend. Accad. Naz. dei Lincei» (8), LIV, 205–209.
- [5] S. S. SINGH (1971–1972) – *On Kählerian recurrent and Ricci-recurrent spaces of second order*, «Atti Accad. delle Scienze di Torino», 106, 509–518.
- [6] S. S. SINGH (1974) – *On Kählerian recurrent spaces of second order with Bochner curvature tensor*, To appear in «Atti Accad. delle Scienze di Torino».
- [7] S. TACHIBANA (1967) – *On the Bochner curvature tensor*, «Nat. Science Report of the Ochanomizu univ.», 18 (1).