

RENDICONTI LINCEI MATEMATICA E APPLICAZIONI

JONATHAN ARAZY, HARALD UPMEIER

Weyl calculus for complex and real symmetric domains

Atti della Accademia Nazionale dei Lincei. Classe di Scienze Fisiche, Matematiche e Naturali. Rendiconti Lincei. Matematica e Applicazioni, Serie 9, Vol. 13 (2002), n.3-4, p. 165–181.

Accademia Nazionale dei Lincei

<http://www.bdim.eu/item?id=RLIN_2002_9_13_3-4_165_0>

L'utilizzo e la stampa di questo documento digitale è consentito liberamente per motivi di ricerca e studio. Non è consentito l'utilizzo dello stesso per motivi commerciali. Tutte le copie di questo documento devono riportare questo avvertimento.

Atti della Accademia Nazionale dei Lincei. Classe di Scienze Fisiche, Matematiche e Naturali. Rendiconti Lincei. Matematica e Applicazioni, Accademia Nazionale dei Lincei, 2002.

JONATHAN ARAZY - HARALD UPMEIER

WEYL CALCULUS FOR COMPLEX AND REAL SYMMETRIC DOMAINS

ABSTRACT. — We define the Weyl functional calculus for real and complex symmetric domains, and compute the associated Weyl transform in the rank 1 case.

KEY WORDS: Functional calculi; Symmetric domains; Weyl transform.

0. INTRODUCTION

In the theory of pseudo-differential operators the Weyl calculus (a quantization method for the cotangent bundle $T^{\sharp}(\mathbb{R}^n)$) is of basic importance since it allows the full symplectic group $\mathrm{Sp}(2n, \mathbb{R})$ as covariance group and the relationship between operators and symbols has optimal continuity properties. Unterberger [10, 11] has introduced an analogous Weyl calculus for (curved) hermitian symmetric spaces of non-compact type and computed the Weyl transform in the simplest case of the unit disk. The higher dimensional case is more difficult. In this paper we define the Weyl calculus for real symmetric domains and then determine the Weyl transform for all symmetric spaces of rank 1. The new feature is the appearance of a hypergeometric function in the spectral decomposition, indicating that the harmonic analysis underlying the Weyl calculus involves (multi-variable) special functions in a significant way.

1. REAL SYMMETRIC DOMAINS AND QUANTIZATION HILBERT SPACES

Real bounded symmetric domains, as defined in [7], are those Riemannian symmetric spaces $D = G/K$ of non-compact type which are real forms of the well-known complex hermitian bounded symmetric domains $D_{\mathbb{C}} = G_{\mathbb{C}}/K_{\mathbb{C}}$, where $G_{\mathbb{C}} = \mathrm{Aut}(D_{\mathbb{C}})^o$ is a real semisimple Lie group of hermitian type and $K_{\mathbb{C}}$ is a maximal compact subgroup. The well-known Harish-Chandra embedding, in its Jordan theoretic form, realizes $D_{\mathbb{C}}$ as the open unit ball

$$D_{\mathbb{C}} = \{z \in Z_{\mathbb{C}} : \|z\| < 1\}$$

of a complex vector space $Z_{\mathbb{C}} \approx \mathbb{C}^n$ endowed with a Jordan triple product $\{uv^*w\}$ [7, 13]. Now let $z \mapsto \bar{z}$ be a conjugation on $Z_{\mathbb{C}}$ preserving the triple product and define

$$Z := \{z \in Z_{\mathbb{C}} : \bar{z} = z\}, \quad D := \{z \in D_{\mathbb{C}} : \bar{z} = z\} = Z \cap D_{\mathbb{C}},$$

$$G := \{g \in G_{\mathbb{C}} : \overline{g(z)} = g(\bar{z}) \quad \forall z \in D_{\mathbb{C}}\} = \{g \in G_{\mathbb{C}} : g(D) = D\}, \quad K := K_{\mathbb{C}} \cap G.$$

Then $D = G/K$ is called a *real bounded symmetric domain* which is a Riemannian symmetric space under the reductive Lie group G . Up to a few low dimensional

exceptions all irreducible Riemannian symmetric spaces of non-compact type can be realized this way. Assuming that Z is irreducible one can show that the following cases occur:

Case 1. $D_{\mathbb{C}}$ is an irreducible hermitian domain with real form D .

We may also include the *flat* case

Case 2. $D_{\mathbb{C}} = Z_{\mathbb{C}} \approx \mathbb{C}^n$, endowed with the usual conjugation, so that $D = Z \approx \mathbb{R}^n$. In this case we obtain (non-reductive) semi-direct products $G_{\mathbb{C}} = U(n) \triangleleft \mathbb{C}^n$, $G = O(n) \triangleleft \mathbb{R}^n$.

Case 3. D is itself a complex hermitian domain with complexification $D_{\mathbb{C}} = D \times \overline{D}$ endowed with the flip conjugation $\overline{(z_1, \overline{z_2})} := (z_2, \overline{z_1})$ for all $z_1, z_2 \in D$. In this case

$$G_{\mathbb{C}} = \{(g_1, \overline{g_2}) : g_1, g_2 \in G\} \approx G \times G$$

where $(g_1, \overline{g_2})(z_1, \overline{z_2}) := (g_1(z_1), \overline{g_2(z_2)})$.

For every real symmetric domain D as above there exists a scale of «quantization Hilbert spaces» H_{ν} of holomorphic functions on the complexification $D_{\mathbb{C}}$ of D . These Hilbert spaces constitute the «scalar holomorphic discrete series» of $G_{\mathbb{C}} = \text{Aut}(D_{\mathbb{C}})^{\circ}$ via irreducible unitary (projective) representations $U_{\nu} : G_{\mathbb{C}} \rightarrow U(H_{\nu})$ of the form

$$(U_{\nu}(g^{-1})h)(z) = j(g, z)h(g(z))$$

for all $g \in G_{\mathbb{C}}$, $h \in H_{\nu}$ and $z \in D_{\mathbb{C}}$. Here $j(g, z)$ is a suitable automorphy factor.

For each irreducible complex bounded symmetric domain B of dimension n , define the *weighted Bergman spaces*

$$H_{\beta}^2(B) := \{h \in L^2(B, d\mu_{\beta}) : h \text{ holomorphic}\}.$$

Here $\beta > p - 1$ is a scalar parameter, where p is the genus of B . The probability measure

$$(1.1) \quad d\mu_{\beta}(z) := \frac{\Gamma_{\Omega_B}(\beta)}{\pi^n \Gamma_{\Omega_B}(\beta - n/r_B)} \Delta(z, z)^{\beta-p} dm(z)$$

involves the so-called Jordan triple determinant $\Delta(z, w)$ and the Gindikin Γ -function of the positive cone Ω_B associated with B . Moreover, r_B is the rank of B and $dm(z)$ is Lebesgue measure. The reproducing kernel of $H_{\beta}^2(B)$ has the form

$$K(z, w) = \Delta(z, w)^{-\beta}$$

for all $z, w \in B$. Returning to the real symmetric domain D with complexification $D_{\mathbb{C}}$, we consider the different cases:

Case 1. If $D_{\mathbb{C}}$ is an irreducible complex symmetric domain with real form D , we define

$$(1.2) \quad H_{\nu} := H_{\nu_{\mathbb{C}}}^2(D_{\mathbb{C}}), \quad \nu_{\mathbb{C}} := 2r\nu/r_{\mathbb{C}}$$

where $r_{\mathbb{C}} \geq r$ is the rank of $D_{\mathbb{C}}$. We have $j(g, z) = (\text{Det } g'(z))^{\nu_{\mathbb{C}}/p_{\mathbb{C}}}$ in this case, where $p_{\mathbb{C}}$ is the genus of $D_{\mathbb{C}}$.

Case 2. In the flat case $Z_{\mathbb{C}} = \mathbb{C}^n$, with real form $Z = \mathbb{R}^n$, the quantization Hilbert spaces are the *Bargmann spaces*

$$H_{\nu} := H_{\nu}^2(\mathbb{C}^n) = \{h \in L^2(\mathbb{C}^n, \mu_{\nu}) : h \text{ holomorphic}\}$$

with respect to the probability measure

$$d\mu_{\nu}(z) = \left(\frac{\nu}{\pi}\right)^n e^{-\nu(z|z)} dm(z)$$

where $(z|w)$ is the scalar product on \mathbb{C}^n and $dm(z)$ is the associated Lebesgue measure. The reproducing kernel is

$$K(z, w) = e^{\nu(z|w)}$$

for all $z, w \in \mathbb{C}^n$ and U_{ν} is the Schrödinger representation of $U(n) \triangleleft \mathbb{C}^n$ in the «complex wave» realization, with its well-known multiplier $j(g, z)$ [1].

Case 3. If D is itself a complex hermitian domain, with measure $d\mu_{\nu}$ as defined in (1.1), we consider the product probability measure $d\mu(z_1, z_2) := d\mu_{\nu}(z_1) d\mu_{\nu}(z_2)$ on $D_{\mathbb{C}} = D \times \overline{D}$ and put

$$H_{\nu} := \{h \in L^2(D_{\mathbb{C}}, d\mu) : h \text{ sesqui-holomorphic}\} = H_{\nu}^2(D) \otimes \overline{H_{\nu}^2(D)}$$

realized via *Hilbert-Schmidt operators*

$$(h\phi)(z) = \int_D d\mu_{\nu}(w) h(z, \overline{w}) \phi(w)$$

for $\phi \in H_{\nu}^2(D)$ and $z \in D$. This Hilbert space has the reproducing kernel

$$(1.3) \quad K(z_1, \overline{z_2}; w_1, \overline{w_2}) = K(z_1, w_1) K(w_2, z_2),$$

with K the kernel function of $H_{\nu}^2(D)$, and

$$U_{\nu}(g_1, \overline{g_2}) h = U_{\nu}(g_1) h U_{\nu}(g_2)^*$$

is the corresponding irreducible unitary (projective) representation of $G_{\mathbb{C}} = G \times G$ on H_{ν} , realized as Hilbert-Schmidt operators. We put $j(g_1, \overline{g_2}; z_1, \overline{z_2}) := j(g_1, z_1) \overline{j(g_2, z_2)}$ in this case.

In all cases the reproducing kernel $K(z, w)$ and the (projective) multiplier $j(g, z)$ are related by

$$(1.4) \quad j(g, z) K(g(z), g(w)) \overline{j(g, w)} = K(z, w)$$

for all $z, w \in D_{\mathbb{C}}$ and $g \in G_{\mathbb{C}}$. This implies

$$(1.5) \quad U_{\nu}(g) K_z = \overline{j(g, z)} K_{g(z)}.$$

2. THE WEYL CALCULUS AND ITS BASIC PROPERTIES

In [1, 2] a general concept of «covariant symbolic calculus» of symmetric domains has been developed. In the (more general) *real* version [2] one considers a linear «symbol»

map $\sigma : H_\nu \rightarrow \{\text{functions on } D\}$ satisfying the covariance condition

$$(2.1) \quad \sigma(U_\nu(g)h) = (\sigma h) \circ g^{-1}$$

for all $g \in G \subset G_{\mathbb{C}}$ and $h \in H_\nu$. More precisely, the domain $\text{Dom}(\sigma)$ should contain all the kernel vectors

$$K_w(z) := K(z, w)$$

for $w \in D_{\mathbb{C}}$, and the condition (2.1) becomes

$$(2.2) \quad \overline{j(g, w)} \sigma(K_{g(w)}) = (\sigma K_w) \circ g^{-1}$$

for all $g \in G$ and $w \in D_{\mathbb{C}}$. In addition we assume that the holomorphic function

$$I(z) := K(z, \bar{z})^{1/2}$$

on $D_{\mathbb{C}}$, defined via the conjugation $z \mapsto \bar{z}$ and the holomorphic square-root on the (simply-connected) domain $D_{\mathbb{C}}$, belongs to $\text{Dom}(\sigma)$ and satisfies

$$(2.3) \quad \sigma I = 1.$$

Since $U_\nu(g)I = I$ for all $g \in G$, σI is a constant function according to (2.1) so that (2.3) is just a normalization.

In [2], the so-called *Toeplitz-Berezin calculus* has been studied in detail (cf. also [3, 9, 15]). We now consider another covariant symbolic calculus, the *Weyl calculus* introduced in the complex setting in [10]. For $\zeta \in D$ the *symmetry* $s_\zeta \in G$ is characterized by the conditions

$$s_\zeta^2 = \text{id}, \quad s_\zeta(\zeta) = \zeta, \quad s'_\zeta(\zeta) = -\text{Id}.$$

LEMMA 2.1. $j(s_\zeta, \zeta) = 1$ for all $\zeta \in D$.

PROOF. Since $s_\zeta^2 = \text{id}$ we have $1 = j(s_\zeta^2, \zeta) = j(s_\zeta, s_\zeta(\zeta)) j(s_\zeta, \zeta) = j(s_\zeta, \zeta)^2$ and hence $j(s_\zeta, \zeta) \in \{\pm 1\}$. Since D is connected it follows that $j(s_\zeta, \zeta) = j(s_0, 0) = 1$. \square

LEMMA 2.2. For $\zeta \in D$ and $z \in D_{\mathbb{C}}$ we have

$$j(s_\zeta, z) = \frac{K(z, \zeta)}{K(s_\zeta z, \zeta)}.$$

PROOF. By Lemma 2.1, we have $j(s_\zeta, z) K(s_\zeta z, \zeta) = j(s_\zeta, z) K(s_\zeta z, s_\zeta \zeta) j(s_\zeta, \zeta) = j(s_\zeta, z) K(s_\zeta z, \zeta) = K(z, \zeta)$. \square

As a special case of Lemma 2.2, we obtain

$$j(s, z) = \frac{K(z, 0)}{K(sz, 0)}$$

for the origin $\zeta = 0 \in D$ and its symmetry $s = s_0$.

DEFINITION 2.1. The *Weyl symbol* map $\omega_\nu : \text{span}\{K_z; z \in D_{\mathbb{C}}\} \rightarrow \mathcal{C}^\infty(D)$ is defined by

$$(2.4) \quad (\omega_\nu K_z)(\zeta) = c_\nu^{-1} \frac{K(\zeta, z)^{1/2}}{K(\zeta, s_\zeta z)^{1/2}} K(\bar{z}, s_\zeta z)^{1/2}$$

for all $z \in D_{\mathbb{C}}$ and $\zeta \in D$. Here \bar{z} is the conjugate of z (so that (2.4) is anti-holomorphic in z). The normalization constant c_{ν} determined by the condition

$$(2.5) \quad \omega_{\nu} I = 1$$

will be computed below. Note that Lemma 2.2 implies

$$(2.6) \quad \overline{(\omega_{\nu} K_z)(\zeta)} = c_{\nu}^{-1} j(s_{\zeta}, z)^{1/2} K(s_{\zeta} z, \bar{z})^{1/2}$$

as a holomorphic function in $z \in D_{\mathbb{C}}$. Together with (1.5), (2.6) implies

$$(2.7) \quad c_{\nu}(\omega_{\nu} K_z)(\zeta) = \overline{(j(s_{\zeta}, z) K(\bar{z}, s_{\zeta} z))^{1/2}} = \overline{(j(s_{\zeta}, z) K_{s_{\zeta} z}(\bar{z}))^{1/2}} = \\ = (U_{\nu}(s_{\zeta}) K_z)(\bar{z})^{1/2} = (K_{\bar{z}} | U_{\nu}(s_{\zeta}) K_z)^{1/2}.$$

EXAMPLE 2.1. In the product case $D_{\mathbb{C}} = D \times \bar{D}$, with D complex hermitian, H_{ν} can be identified with the space of Hilbert-Schmidt operators acting on $H_{\nu}^2(D)$, and under this identification

$$K_{z_1, \bar{z}_2} = K_{z_1} K_{z_2}^* \quad (z_1, z_2 \in D)$$

becomes a rank 1 operator [2, Example 3.1]. Therefore (2.7) yields

$$c_{\nu}(\omega_{\nu} K_{z_1, \bar{z}_2})(\zeta) = (K_{z_2, \bar{z}_1} | U_{\nu}(s_{\zeta}) K_{z_1, \bar{z}_2})^{1/2} = (K_{z_2} K_{z_1}^* | U_{\nu}(s_{\zeta}) (K_{z_1} K_{z_2}^*))^{1/2} = \\ = (K_{z_2} K_{z_1}^* | U_{\nu}(s_{\zeta}) K_{z_1} K_{z_2}^* U_{\nu}(s_{\zeta})^*)^{1/2} = (K_{z_2} K_{z_1}^* | (U_{\nu}(s_{\zeta}) K_{z_1})(U_{\nu}(s_{\zeta}) K_{z_2})^*)^{1/2} = \\ = [(K_{z_2} | U_{\nu}(s_{\zeta}) K_{z_1})(U_{\nu}(s_{\zeta}) K_{z_2} | K_{z_1})]^{1/2} = (K_{z_2} | U_{\nu}(s_{\zeta}) K_{z_1}) = \text{tr}(U_{\nu}(s_{\zeta}) K_{z_1} K_{z_2}^*).$$

Hence we have

$$c_{\nu}(\omega_{\nu} T)(\zeta) = \text{tr}(U_{\nu}(s_{\zeta}) T)$$

for all (trace-class) operators T acting on $H_{\nu}^2(D)$. This coincides with the «Weyl symbol» of T as defined in [10].

PROPOSITION 2.1. *The Weyl symbol (2.4) is covariant under G .*

PROOF. Let $\zeta \in D$, $g \in G$ and $z \in D_{\mathbb{C}}$. Then $s_{g(\zeta)} = g s_{\zeta} g^{-1}$ and (1.2) and (2.6) imply

$$c_{\nu} [j(g, z) \overline{(\omega_{\nu} K_{g(z)})(g(\zeta))}]^2 = j(g, z)^2 j(s_{g(\zeta)}, g(z)) K(s_{g(\zeta)}(g(z)), \overline{g(\zeta)}) = \\ = j(g, z) j(g s_{\zeta} g^{-1}, g(z)) K(g(s_{\zeta}(z)), g(\bar{z})) \overline{j(g, \bar{z})} = \\ = j(s_{\zeta}, z) j(g, s_{\zeta}(z)) K(g(s_{\zeta}(z)), g(\bar{z})) \overline{j(g, \bar{z})} = \\ = j(s_{\zeta}, z) K(s_{\zeta}(z), \bar{z}) = c_{\nu} \overline{(\omega_{\nu} K_z)(\zeta)}^2$$

since $j(g, z) j(g s_{\zeta} g^{-1}, g(z)) = j(g s_{\zeta}, z) = j(s_{\zeta}, z) j(g, s_{\zeta}(z))$. Taking holomorphic square-roots and conjugates it follows that $\overline{j(g, \bar{z})} (\omega_{\nu} K_{g(z)})(g(\zeta)) = (\omega_{\nu} K_z)(\zeta)$ which yields covariance in view of (2.2). \square

Given a covariant symbolic calculus σ one defines its *adjoint* $\sigma^* : \{\text{functions on } D\} \rightarrow \rightarrow H_\nu$ by assigning to a function $f \in \text{Dom}(\sigma^*)$ the holomorphic function

$$(\sigma^* f)(z) := \int_D d\mu_0(\zeta) f(\zeta) \overline{(\sigma K_z)(\zeta)}$$

on $D_{\mathbb{C}}$. Here $d\mu_0$ is the G -invariant measure on D normalized by the condition

$$(2.8) \quad \int_D d\mu_0(\zeta) I(\zeta)^{-1} = 1.$$

By [2, Proposition 4.4] σ^* is the adjoint of σ with respect to $L^2(D, d\mu_0)$. According to Definition 2.1 the adjoint $f \mapsto \omega_\nu^* f$ of the Weyl symbol map is given by

$$\begin{aligned} (\omega_\nu^* f)(z) &= \int_D d\mu_0(\zeta) f(\zeta) \overline{(\omega_\nu K_z)(\zeta)} = c_\nu^{-1} \int_D d\mu_0(\zeta) f(\zeta) \frac{K(z, \zeta)^{1/2}}{K(s_\zeta z, \zeta)^{1/2}} K(s_\zeta z, \bar{z})^{1/2} = \\ &= c_\nu^{-1} \int_D d\mu_0(\zeta) f(\zeta) j(s_\zeta, z)^{1/2} K(s_\zeta z, \bar{z})^{1/2} = c_\nu^{-1} \int_D d\mu_0(\zeta) f(\zeta) \overline{(U_\nu(s_\zeta) K_z)(\bar{z})}^{1/2} \end{aligned}$$

as a holomorphic function in $z \in D_{\mathbb{C}}$. Whereas the Toeplitz map $f \mapsto \tau_\nu^* f$ is well-defined for $f \in L^\infty(D)$, it is more difficult to find conditions on $f \in C^\infty(D)$ such that $\omega_\nu^* f$ is well-behaved [11, 14].

EXAMPLE 2.2. In the flat case $D = \mathbb{R}^n$ and $D_{\mathbb{C}} = \mathbb{C}^n$, the ν -th Bargmann space $H_\nu^2(\mathbb{C}^n)$ has the reproducing kernel $K(z, w) = \exp \nu(z|w)$ and

$$s_\zeta z = 2\zeta - z$$

is the symmetry. Hence the Weyl calculus $\omega_\nu^* : L^2(\mathbb{R}^n) \rightarrow H_\nu^2(\mathbb{C}^n)$ has the form

$$\begin{aligned} c_\nu (\omega_\nu^* f)(z) &= \int_{\mathbb{R}^n} d\zeta f(\zeta) \cdot \exp \frac{\nu}{2} ((z|\zeta) + (s_\zeta z|\bar{z}) - (s_\zeta z|\zeta)) = \\ &= \int_{\mathbb{R}^n} d\zeta f(\zeta) \exp \frac{\nu}{2} ((z|\zeta) + (2\zeta - z|\bar{z} - \zeta)) = \\ &= \int_{\mathbb{R}^n} d\zeta f(\zeta) \exp \left(2\nu(z|\zeta) - \frac{\nu}{2}(z|\bar{z}) - \nu(\zeta|\zeta) \right). \end{aligned}$$

Since $c_\nu = 2^{-n/4}$ in this case, we obtain the *Bargmann transform* [5, p. 40].

The link *transform* of a covariant symbolic calculus σ is defined as the map $f \mapsto \mapsto (\sigma \sigma^*) f := \sigma(\sigma^* f)$ acting on functions on D . By [2, Proposition 4.7] $\sigma \sigma^*$ is an integral operator in $L^2(D, d\mu_0)$ with kernel

$$\int_{D_{\mathbb{C}}} d\mu_\nu(z) (\sigma K_z)(\xi) \overline{(\sigma K_z)(\eta)}$$

for $\xi, \eta \in D$. For the *Weyl transform* $\omega_\nu \omega_\nu^*$ we obtain the integral kernel

$$c_\nu^{-2} \int_{D_{\mathbb{C}}} d\mu_\nu(z) \left(\frac{K(\xi, z) K(\bar{z}, s_\zeta z) K(z, \eta) K(s_\eta z, \bar{z})}{K(\xi, s_\zeta z) K(s_\eta z, \eta)} \right)^{1/2}$$

for $\xi, \eta \in D$. In comparison, the integral kernel of the *Toeplitz transform* $\tau_\nu \tau_\nu^*$ has the much simpler expression [2, (5.4)]

$$\frac{K(\xi, \eta)}{K(\xi, \xi)^{1/2} K(\eta, \eta)^{1/2}}.$$

Since $D = G/K$ is a Riemannian symmetric space there is an explicit *Plancherel decomposition* [6]

$$L^2(D, \mu_0) = \int_{\mathfrak{a}^\sharp} d\lambda |c(\lambda)|^{-2} \langle G \rangle_\lambda$$

in terms of the Hilbert spaces $\langle G \rangle_\lambda$ of the principal series of G with parameter $\lambda \in \mathfrak{a}^\sharp$, where $c(\lambda)$ is Harish-Chandra's c -function and we use an Iwasawa decomposition $G = NAK$ with $\mathfrak{a} := \text{Lie}(A)$. By covariance every covariant symbolic calculus σ yields a multiplicity-free decomposition

$$H_\nu \approx \int d\sigma_0(\lambda) \langle G \rangle_\lambda$$

of H_ν , under the restricted action of $G \subset G_{\mathbb{C}}$. The defining measure $d\sigma_0(\lambda)$ depends on the choice of calculus, more precisely on the *eigenvalues*

$$\widetilde{\sigma\sigma^*}(\lambda) := (\sigma\sigma^*\phi_\lambda)(0)$$

of the G -invariant link transform $\sigma\sigma^*$, computed on the spherical function ϕ_λ of type $\lambda \in \mathfrak{a}^\sharp$. Here $0 \in D$ is the origin. It is technically easier to use the NA -covariant «exponential functions» e_λ , where

$$\phi_\lambda(\zeta) = \int_K dk e_\lambda(k\zeta)$$

for all $\zeta \in D$. For the Toeplitz-Berezin calculus τ_ν the eigenvalues of $\tau_\nu \tau_\nu^*$ are given by the integral

$$\widetilde{\tau_\nu \tau_\nu^*}(\lambda) = (\tau_\nu^* e_\lambda)(0) = \int_D d\mu_0(\zeta) e_\lambda(\zeta) (\tau_\nu K_0)(\zeta) = \int_D d\mu_0(\zeta) e_\lambda(\zeta) K(0, \zeta) K(\zeta, \zeta)^{-1/2}$$

which can be computed using the structure theory of Jordan triples [2, 15, 3, 9] yielding a (complicated) product of classical Γ -functions. For arbitrary covariant symbolic calculi σ_1, σ_2 on H_ν there is a «product formula» [2, Theorem 4.9]

$$\widetilde{\sigma_1 \sigma_2^*}(\lambda) = \frac{(\sigma_1^* e_\lambda)(0) (\sigma_2^* e_\lambda)(0)}{(\tau_\nu^* e_\lambda)(0)}.$$

Thus the integral

$$(\omega_\nu^* e_\lambda)(0) = \int_D d\mu_0(\zeta) e_\lambda(\zeta) (\omega_\nu K_0)(\zeta) = c_\nu^{-1} \int_D d\mu_0(\zeta) e_\lambda(\zeta) \frac{K(0, \zeta)^{1/2}}{K(\zeta, s_\zeta(0))^{1/2}} K(0, s_\zeta(0))^{1/2}$$

is needed for the computation of the eigenvalues of $\widetilde{\omega_\nu \omega_\nu^*}(\lambda)$ of the Weyl transform.

3. POLAR COORDINATES AND ROOT DECOMPOSITION

From now on we only consider the non-flat case. For a deeper study of the Weyl calculus (and other covariant symbolic calculi) on real symmetric domains it is necessary to recall the basic structure theory of symmetric spaces G/K (of non-compact type) related to the *root decomposition* [6]

$$\mathfrak{g} = \mathfrak{a} \oplus \mathfrak{m} \oplus \int_{\alpha \in \Sigma} \mathfrak{g}_\alpha$$

induced by a Cartan subspace $\mathfrak{a} \subset \mathfrak{g}$. Here \mathfrak{m} is the centralizer of \mathfrak{a} in \mathfrak{k} and \mathfrak{g}_α denotes the root space associated with $\alpha \in \mathfrak{a}^\#$. Put $\Sigma := \{\alpha \in \mathfrak{a}^\# \setminus \{0\} : \mathfrak{g}_\alpha \neq \{0\}\}$. According to [7] every (irreducible) real symmetric domain $D = G/K$ has an unbounded realization as a *real Siegel domain*

$$D \approx \{x + y + v \in X \oplus Y \oplus V : x - \{e v^* v\} \in \Omega\}.$$

Here X is a euclidean Jordan algebra of rank r , with unit element e and positive cone Ω , $X \oplus Y$ is a semi-simple real Jordan $*$ -algebra with self-adjoint part X and skew-adjoint part Y , and the Peirce decomposition [7] of Z with respect to e has the 1-eigenspace $X \oplus Y$ and the $\frac{1}{2}$ -eigenspace V . Now choose a *frame* e_1, \dots, e_r of minimal idempotents in X satisfying $e_1 + \dots + e_r = e$ and consider the joint Peirce decomposition [7]

$$X = \sum_{1 \leq i \leq j \leq r} X_{ij}, \quad Y = \sum_{1 \leq i \leq j \leq r} Y_{ij}, \quad V = \sum_{1 \leq j \leq r} V_{0j}.$$

Then for $1 \leq j \leq r$ and $1 \leq i < j \leq r$ we have $\dim X_{ij} = 1$ and

$$a := \dim X_{ij}, \quad b := \dim V_{0j}, \quad c := \dim Y_{ij}$$

are independent of i, j and of the frame e_1, \dots, e_r . For the symmetric cones we have $Y = \{0\} = V$. In all other cases (except root system D_2 , which is not considered in the sequel) the classification yields $\dim Y_{ij} = a \quad (1 \leq i < j \leq r)$. Hence the fine structure of D is completely encoded in the numerical invariants a, b, c . In particular, we have $n_X := \dim X = r + \frac{a}{2} r(r-1)$, $n_Y := \dim Y = cr + \frac{a}{2} r(r-1)$, $n_V := \dim V = br$. Returning to the bounded realization, the commuting completely integrable holomorphic vector fields

$$(3.1) \quad M_j := (e_j - \{z e_j^* z\}) \frac{\partial}{\partial z} \quad (1 \leq j \leq r)$$

on $D_{\mathbb{C}}$ [13, 7] leave D invariant and can be chosen as a basis of \mathfrak{a} . Let $M_1^\#, \dots, M_r^\# \in \mathfrak{a}^\#$ denote the dual basis satisfying $M_i^\#(M_j) = \delta_{ij}$ for $1 \leq i, j \leq r$. Then, by [7, 15] the positive restricted roots of \mathfrak{g} are the following:

$$(3.2) \quad M_j^\# - M_i^\#, \quad \text{multiplicity } a \quad (1 \leq i < j \leq r)$$

$$(3.3) \quad M_j^\# + M_i^\#, \quad \text{multiplicity } a \quad (1 \leq i < j \leq r)$$

$$(3.4) \quad 2M_j^\#, \quad \text{multiplicity } c \quad (1 \leq j \leq r)$$

$$(3.5) \quad M_j^\sharp, \quad \text{multiplicity } b \quad (1 \leq j \leq r),$$

unless $Z = X$, in which case only (3.2) occurs. This case and also the root type D_2 will be omitted in the sequel. For the half-sum of positive roots ρ we obtain

$$\begin{aligned} 2\rho &= \sum_{\alpha \in \Sigma_+} m_\alpha \alpha = \sum_{1 \leq i < j \leq r} a (M_j^\sharp - M_i^\sharp + M_j^\sharp + M_i^\sharp) + \sum_{1 \leq j \leq r} (c 2M_j^\sharp + b M_j^\sharp) = \\ &= \sum_{1 \leq i < j \leq r} 2a M_j^\sharp + \sum_{1 \leq j \leq r} (2c + b) M_j^\sharp = \sum_{1 \leq j \leq r} 2 \left((j-1)a + c + \frac{b}{2} \right) M_j^\sharp \end{aligned}$$

and hence

$$\rho = \sum_{1 \leq j \leq r} \left((j-1)a + c + \frac{b}{2} \right) M_j^\sharp.$$

By [6, Theorem 5.8] there exists a Haar measure dg on G such that

$$\int_G dg f(g(0)) = \int_{\mathbb{R}_+^r} dt_1 \cdots dt_r f(\exp(\sum_j t_j M_j)(0)) \prod_{\alpha \in \Sigma_+} \sinh(\alpha(\sum_j t_j M_j))^{m_\alpha}$$

holds for K -invariant functions of f on $D = G/K$. Here Σ_+ denotes the set of positive roots and m_α is the multiplicity of $\alpha \in \Sigma_+$. For each tripotent $c = \{c c^* c\} \in Z$ the vector field

$$M_c := (c - \{z c^* z\}) \frac{\partial}{\partial z}$$

satisfies $\exp(t M_c)(0) = \tanh(t) c$ for all $t \in \mathbb{R}$ [13, 7]. Similarly, we have

$$\exp\left(\sum_{j=1}^r t_j M_j\right)(0) = \sum_{j=1}^r \tanh(t_j) e_j$$

for the (commuting) vector fields (3.1). Using the coordinates

$$(3.6) \quad x_j = \tanh^2(t_j) \in [0, 1]$$

satisfying $\frac{dx_j}{dt_j} = 2x_j^{1/2}(1-x_j)$, the explicit root decomposition (3.2)-(3.5) yields

$$\begin{aligned} &\int_G dg f(g(0)) = \\ &= \int_{\mathbb{R}_+^r} \prod_j dt_j \sinh(2t_j)^c \sinh(t_j)^b \cdot f(\sum_j \tanh(t_j) e_j) \cdot \prod_{i < j} |\sinh(t_j - t_i) \sinh(t_j + t_i)|^a = \\ &= \int_{[0,1]^r} \prod_j \frac{dx_j}{2(1-x_j)x_j^{1/2}} \left(\frac{2x_j^{1/2}}{1-x_j}\right)^c \left(\frac{x_j}{1-x_j}\right)^{b/2} \cdot \prod_{i < j} \left|\frac{x_i - x_j}{(1-x_i)(1-x_j)}\right|^a \cdot f(\sum_j x_j^{1/2} e_j) = \\ &= 2^{r(c-1)} \int_{[0,1]^r} dx_1 \cdots dx_r \cdot \prod_j (1-x_j)^{-1-c-b/2-a(r-1)} x_j^{(c-1+b)/2} \cdot \prod_{i < j} |x_i - x_j|^a \cdot f(\sum_j x_j^{1/2} e_j) = \\ &= + 2^{n_Y - n_X} \int_{[0,1]^r} f(\sum_j x_j^{1/2} e_j) \prod_{i < j} |x_i - x_j|^a \cdot \prod_j dx_j (1-x_j)^{-(n_X + n_Y + n_V/2)/r} x_j^{(n_Y - n_X + n_V)/2r}. \end{aligned}$$

Let Ω be the positive cone of the euclidean Jordan algebra X . The Gindikin Γ -function Γ_Ω associated with Ω [4, Chapter VII] has the property [4, pp. 123 and 104]

$$(3.7) \quad \Gamma_\Omega\left(\alpha + \frac{n_X}{r}\right) = c_\Omega \int_{\mathbb{R}_+^r} \prod_{i < j} |x_i - x_j|^a \prod_j dx_j e^{-x_j} x_j^\alpha,$$

where c_Ω is a constant depending only on Ω . Similarly for the Beta-integral [4, pp. 130 and 104] which is symmetric in α and γ :

$$(3.8) \quad \frac{\Gamma_\Omega\left(\alpha + \frac{n_X}{r}\right) \Gamma_\Omega\left(\gamma + \frac{n_X}{r}\right)}{\Gamma_\Omega\left(\alpha + \gamma + \frac{2n_X}{r}\right)} = c_\Omega \int_{[0,1]^r} \prod_{i < j} |x_i - x_j|^a \prod_j dx_j x_j^\alpha (1 - x_j)^\gamma.$$

LEMMA 3.1. *For $\xi, \eta \in D \cap X$ we have*

$$K(\xi, \eta)^{-1/2} = \Delta(e - \{\xi e^* \eta\})^\nu.$$

PROOF. In case $D_{\mathbb{C}}$ is irreducible, we have

$$K(\xi, \eta)^{-1/2} = \Delta_{\mathbb{C}}(\xi, \eta)^{\nu_{\mathbb{C}}/2} = \Delta_{\mathbb{C}}(e - \{\xi e^* \eta\})^{\nu_{\mathbb{C}}/2},$$

$$\Delta(e - \{\xi e^* \eta\})^{\nu_{\mathbb{C}} \nu_{\mathbb{C}}/2r} = \Delta(e - \{\xi e^* \eta\})^\nu.$$

In case $D_{\mathbb{C}} = D \times \overline{D}$, with D complex hermitian, (1.3) implies

$$K(\xi, \xi, \eta, \eta)^{-1/2} = K(\xi, \eta)^{-1} = \Delta(\xi, \eta)^\nu = \Delta(e - \{\xi e^* \eta\})^\nu.$$

In both cases, the assertion follows. \square

PROPOSITION 3.1. *The measure μ_0 normalized by (2.8) is given by*

$$\int_D d\mu_0(\zeta) f(\zeta) = c_\Omega 2^{n_X - n_Y} \frac{\Gamma_\Omega\left(\nu + \frac{n_X - n_Y}{2r}\right)}{\Gamma_\Omega\left(\nu - \frac{n_Y + n_V/2}{r}\right) \Gamma_\Omega\left(\frac{n}{2r}\right)} \int_G dg f(g(0)).$$

PROOF. By Lemma 3.1 we have for $\zeta := \sum_j \tanh(t_j) e_j \in D \cap X$

$$I(\zeta)^{-1} = K(\zeta, \zeta)^{-1/2} = \Delta(e - \{\zeta e^* \zeta\})^\nu = \prod_j (1 - \tanh^2(t_j))^\nu = \prod_j (1 - x_j)^\nu.$$

Since $n - 2n_Y - n_V = n_X - n_Y$, it follows that

$$\begin{aligned} \int_G dg I(g(0))^{-1} &= 2^{n_Y - n_X} \int_{[0,1]^r} \prod_{i < j} |x_i - x_j|^a \cdot \prod_j dx_j (1 - x_j)^{\nu - (n_X + n_Y + n_V/2)/r} x_j^{(n_Y - n_X + n_V)/2r} = \\ &= c_\Omega^{-1} 2^{n_Y - n_X} \frac{\Gamma_\Omega\left(\nu - \frac{n_Y + n_V/2}{r}\right) \Gamma_\Omega\left(\frac{n}{2r}\right)}{\Gamma_\Omega\left(\nu + \frac{n_X - n_Y}{2r}\right)}. \quad \square \end{aligned}$$

The spherical functions of the cone Ω , regarded as a reductive symmetric space, can be expressed in terms of the so-called *Jack polynomials* $J_{\underline{m}}(x_1, \dots, x_r)$ associated with an integer partition $\underline{m} = (m_1, \dots, m_r)$. Using the Jack polynomials and the multi-variable *Pochhammer symbol*

$$(\alpha)_{\underline{m}} := \frac{\Gamma_{\Omega}(\alpha + \underline{m})}{\Gamma_{\Omega}(\alpha)}$$

one defines the multivariable *hypergeometric series* [4]

$${}_2F_1\left(\begin{matrix} \alpha & \beta \\ \gamma \end{matrix} \right) (x_1, \dots, x_r) = \sum_{\underline{m}} \frac{(\alpha)_{\underline{m}} (\beta)_{\underline{m}}}{(\gamma)_{\underline{m}} (1)_{\underline{m}}} J_{\underline{m}}(x_1, \dots, x_r).$$

The multivariable hypergeometric function together with the Gindikin Γ -function yields the following Selberg-type integral

$$\begin{aligned} c_{\Omega} \cdot \int_{[0,1]^r} \prod_j dz_j z_j^{\alpha} (1 - z_j)^{\gamma} \left(1 - \frac{z_j}{2}\right)^{-\beta} \prod_{i < j} |z_i - z_j|^a = \\ (3.9) \quad = \frac{\Gamma_{\Omega}\left(\alpha + \frac{n_X}{r}\right) \Gamma_{\Omega}\left(\gamma + \frac{n_X}{r}\right)}{\Gamma_{\Omega}\left(\alpha + \gamma + \frac{2n_X}{r}\right)} {}_2F_1\left(\begin{matrix} \alpha + \frac{n_X}{r} & \beta \\ \alpha + \gamma + \frac{2n_X}{r} \end{matrix} \right) \left(\frac{1}{2}, \dots, \frac{1}{2}\right) \end{aligned}$$

for (suitably restricted) scalar parameters α, β, γ . Since $K(z, 0) = 1$ in the bounded setting, (2.4) implies

$$(3.10) \quad (\omega_{\nu} K_0)(\zeta) = c_{\nu}^{-1} \mathfrak{b}_{\nu}(\zeta),$$

where we define $\mathfrak{b}_{\nu}(\zeta) := K(s_{\zeta}(0), \zeta)^{-1/2}$ for all $\zeta \in D$.

LEMMA 3.2. *The K -invariant function \mathfrak{b}_{ν} on D satisfies*

$$\mathfrak{b}_{\nu} \left(\sum_{j=1}^r \tanh(t_j) e_j \right) = \prod_{j=1}^r \left[\frac{1 - \tanh^2(t_j)}{1 + \tanh^2(t_j)} \right]^{\nu} = \prod_{j=1}^r \left(\frac{1 - x_j}{1 + x_j} \right)^{\nu}.$$

PROOF. By Lemma 3.1 we have $K(\zeta, s_{\zeta}(0))^{-1/2} = \Delta(e - \{\zeta e^* s_{\zeta}(0)\})^{\nu}$ for $\zeta \in D \cap X$. Applying geodesic reflection, it follows that for $\xi := \sum_{j=1}^r \tanh(t_j) e_j \in D \cap X$ we have

$$s_{\xi}(0) = \sum_{j=1}^r \tanh(2t_j) e_j.$$

By orthogonality of $\{e_j\}$, we obtain

$$\begin{aligned} \mathfrak{b}_\nu \left(\sum_{j=1}^r \tanh(t_j) e_j \right) &= \Delta \left(e - \left\{ \left(\sum_{j=1}^r \tanh(t_j) e_j \right) e^* \left(\sum_{k=1}^r \tanh(2t_k) e_k \right) \right\} \right)^\nu = \\ &= \prod_{j=1}^r (1 - \tanh(t_j) \tanh(2t_j))^\nu = \\ &= \prod_{j=1}^r \left(1 - 2 \frac{\tanh^2(t_j)}{1 + \tanh^2(t_j)} \right)^\nu = \prod_{j=1}^r \left(\frac{1 - \tanh^2(t_j)}{1 + \tanh^2(t_j)} \right)^\nu. \quad \square \end{aligned}$$

PROPOSITION 3.2. *The normalizing constant for the Weyl calculus at parameter ν is*

$$c_\nu = 2^{-\nu} {}_2F_1 \left(\nu - \frac{n_Y + n_V/2}{\nu + \frac{n_X - n_Y}{2r}} \quad \nu \right) \left(\frac{1}{2}, \dots, \frac{1}{2} \right).$$

PROOF. By definition, c_ν is chosen such that $\omega_\nu I = 1$. Since the definition of ω_ν applies directly only to the kernel vectors we use the dual condition $\omega_\nu^* 1 = I$. By (3.10) we have

$$1 = (\omega_\nu^* 1)(0) = \int_D d\mu_0(\zeta) (\omega_\nu K_0)(\zeta) = c_\nu^{-1} \int_D d\mu_0(\zeta) \mathfrak{b}_\nu(\zeta)$$

and hence

$$c_\nu = \int_D d\mu_0(\zeta) \mathfrak{b}_\nu(\zeta).$$

Applying Proposition 3.1 and Lemma 3.2 yields, putting $z_j = 1 - x_j$,

$$\begin{aligned} &\frac{\Gamma_\Omega \left(\nu - \frac{n_Y + n_V/2}{r} \right) \Gamma_\Omega \left(\frac{n}{2r} \right)}{\Gamma_\Omega \left(\nu + \frac{n_X - n_Y}{2r} \right)} \int_D d\mu_0(\zeta) \mathfrak{b}_\nu(\zeta) = c_\Omega 2^{n_X - n_Y} \int_G dg \mathfrak{b}_\nu(g(0)) = \\ &= c_\Omega \int_{[0,1]^r} \prod_{i < j} |z_i - z_j|^a \cdot \prod_j dz_j \cdot z_j^{-(n_X + n_Y + n_V/2)/r} (1 - z_j)^{(n_Y - n_X + n_V)/2r} \left(\frac{z_j}{2 - z_j} \right)^\nu = \\ &= 2^{-\nu} c_\Omega \int_{[0,1]^r} \prod_{i < j} |z_i - z_j|^a \cdot \prod_j dz_j \cdot z_j^{\nu - (n_X + n_Y + n_V/2)/r} (1 - z_j)^{(n_Y - n_X + n_V)/2r} \left(1 - \frac{z_j}{2} \right)^{-\nu} = \\ &= 2^{-\nu} \frac{\Gamma_\Omega \left(\nu - \frac{n_Y + n_V/2}{r} \right) \Gamma_\Omega \left(\frac{n}{2r} \right)}{\Gamma_\Omega \left(\nu + \frac{n_X - n_Y}{2r} \right)} {}_2F_1 \left(\nu - \frac{n_Y + n_V/2}{\nu + \frac{n_X - n_Y}{2r}} \quad \nu \right) \left(\frac{1}{2}, \dots, \frac{1}{2} \right). \quad \square \end{aligned}$$

In the sequel the *Laplace-Beltrami operator* Δ on D (not to be confused with the Jordan determinant) will play a crucial role.

PROPOSITION 3.3. *Expressed in the coordinates*

$$(3.11) \quad y_j := -\sinh^2(t_j) \quad (1 \leq j \leq r)$$

the Laplace-Beltrami operator Δ on D has the K -radial part $\tilde{\Delta}$ given by

$$-\frac{1}{4} \tilde{\Delta} = \sum_{j=1}^r \left\{ y_j(1-y_j) \frac{\partial}{\partial y_j} + a \sum_{i \neq j} \frac{y_j(1-y_j)}{y_i-y_j} + \frac{1+c+b}{2} - \left(1+c+\frac{b}{2}\right) y_j \right\} \frac{\partial}{\partial y_j}.$$

PROOF. For any Riemannian symmetric space G/K , the K -radial part of Δ realized on $A_+ := \exp(\mathfrak{a}_+)(0)$ has the form [6, Proposition II. 3.9]

$$\tilde{\Delta} = \Delta_A + \sum_{\alpha \in \Sigma_+} m_\alpha \coth(\alpha) \alpha_\sharp$$

where Δ_A is the (euclidean) Laplacian on A and $\alpha_\sharp \in \mathfrak{a}$ is determined by $\langle \alpha_\sharp | H \rangle = \alpha(H)$ for all $H \in \mathfrak{a}$. Specializing to the root decomposition (3.2)-(3.5) we obtain

$$\begin{aligned} \tilde{\Delta} &= \sum_j \frac{\partial^2}{\partial t_j^2} + \sum_j \left(c \coth(2t_j) 2 \frac{\partial}{\partial t_j} + b \coth(t_j) \frac{\partial}{\partial t_j} \right) + \\ &+ a \sum_{i < j} \coth(t_j - t_i) \left(\frac{\partial}{\partial t_j} - \frac{\partial}{\partial t_i} \right) + \coth(t_j + t_i) \left(\frac{\partial}{\partial t_j} + \frac{\partial}{\partial t_i} \right) = \\ &= \sum_j \left[\frac{\partial}{\partial t_j} + 2c \coth(2t_j) + b \coth(t_j) + a \sum_{i \neq j} (\coth(t_j - t_i) + \coth(t_j + t_i)) \right] \frac{\partial}{\partial t_j} \end{aligned}$$

since $-\coth(t_j - t_i) = \coth(t_i - t_j)$. Since $\frac{dy_j}{dt_j} = -\sinh^2(t_j)$ we have

$$\coth(2t_j) \frac{\partial}{\partial t_j} = -\coth(2t_j) \sinh(2t_j) \frac{\partial}{\partial y_j} = (2y_j - 1) \frac{\partial}{\partial y_j},$$

$$\coth(t_j) \frac{\partial}{\partial t_j} = -\coth(t_j) \sinh(2t_j) \frac{\partial}{\partial y_j} = 2(y_j - 1) \frac{\partial}{\partial y_j},$$

$$(\coth(t_j - t_i) + \coth(t_j + t_i)) \frac{\partial}{\partial t_j} = \frac{\sinh(2t_j) \sinh(2t_i)}{\sinh(t_j - t_i) \sinh(t_j + t_i)} \frac{\partial}{\partial y_j},$$

$$\frac{4 \cosh^2(t_j) \sinh^2(t_j)}{\sinh^2(t_j) \cosh^2(t_i) - \cosh^2(t_j) \sinh^2(t_i)} \frac{\partial}{\partial y_j} = -\frac{4y_j(1-y_j)}{y_i-y_j} \frac{\partial}{\partial y_j}$$

and

$$\left(\frac{\partial}{\partial t_j}\right)^2 = \left(\frac{dy_j}{dt_j}\right)^2 \left(\frac{\partial}{\partial y_j}\right)^2 + \frac{d^2 y_j}{dt_j^2} \frac{\partial}{\partial y_j} = -4y_j(1-y_j) \left(\frac{\partial}{\partial y_j}\right)^2 + (4y_j - 2) \frac{\partial}{\partial y_j}.$$

Substituting into the previous expression for $\tilde{\Delta}$ yields the assertion. \square

4. THE WEYL TRANSFORM FOR RANK 1 DOMAINS

The Weyl transform is harder to analyze than the Toeplitz-Berezin transform. Up to now only the simplest case of the unit disk has been treated in detail [10]. In this section we analyze the Weyl transform for an important class of higher-dimensional symmetric domains, namely those of rank 1. This includes the unit ball in \mathbb{C}^n , and our main result is new even in this special case.

Let \mathbb{K} denote one of the real division algebras $\mathbb{R}, \mathbb{C}, \mathbb{H}$ or \mathbb{O} . Then $Z := \mathbb{K}^m$ becomes a real Jordan triple for $m \geq 1$, with $m = 2$ in case $\mathbb{K} = \mathbb{O}$. The unit balls

$$D := \left\{ (x_1, \dots, x_m) \in \mathbb{K}^m : \sum_{i=1}^m x_i x_i^* < 1 \right\}$$

are precisely the real bounded symmetric domains of rank 1. Here $x \mapsto x^*$ is the natural involution on \mathbb{K} . We put

$$a := \dim_{\mathbb{R}} \mathbb{K}, \quad n := \dim_{\mathbb{R}} D = am.$$

The Peirce decomposition with respect to the tripotent $e := (1, 0, \dots, 0) \in \mathbb{K}^m$ has the form

$$\mathbb{K}^m = X \oplus Y \oplus V$$

where

$$X = \mathbb{R} \times \{0\}^{m-1}, \quad Y = \mathbb{R}^\perp \times \{0\}^{m-1}, \quad V = \{0\} \times \mathbb{K}^{m-1}$$

and $\mathbb{R}^\perp := \{x \in \mathbb{K} : x^* = -x\}$. The vector field

$$M := (e - \{z e^* z\}) \frac{\partial}{\partial z}$$

generating \mathfrak{a} gives rise to the positive roots

$$(4.1) \quad 2M^\sharp, \quad \text{multiplicity } c = a - 1$$

$$(4.2) \quad M^\sharp, \quad \text{multiplicity } b = n - a.$$

For the half-sum ρ of positive roots we obtain

$$2\rho = (a - 1)2M^\sharp + (n - a)M^\sharp = \left(-1 + \frac{n + a}{2}\right) 2M^\sharp.$$

In terms of the coordinate $y := -\sinh^2(t) \in (-\infty, 0]$ the Laplace-Beltrami operator Δ

on D has the radial part (Proposition 3.2)

$$(4.3) \quad -\frac{\tilde{\Delta}}{4} = y(1-y) \left(\frac{d}{dy}\right)^2 + \left(\frac{n}{2} - \frac{n+a}{2}y\right) \frac{d}{dy}$$

when acting on K -invariant functions. Since (4.3) corresponds to the hypergeometric equation it follows that the hypergeometric series

$${}_2F_1\left(\begin{matrix} \alpha & \beta \\ \gamma \end{matrix}\right)(y) := \sum_{k \geq 0} \frac{(\alpha)_k (\beta)_k}{(\gamma)_k} \frac{y^k}{k!},$$

yields the eigenfunctions (regular at $y = 0$):

$$-\frac{\tilde{\Delta}}{4} {}_2F_1\left(\begin{matrix} \alpha & \beta \\ n/2 \end{matrix}\right) = \alpha\beta {}_2F_1\left(\begin{matrix} \alpha & \beta \\ n/2 \end{matrix}\right)$$

where

$$(4.4) \quad \alpha + \beta = \frac{n+a}{2} - 1 = 2\rho.$$

Using the coordinate $x := \tanh^2(t)$ we obtain the *spherical function*

$$(4.5) \quad \begin{aligned} \phi_\lambda(x^{1/2}e) &= \phi_\lambda(\tanh(t)e) = {}_2F_1\left(\begin{matrix} \rho+\lambda & \rho-\lambda \\ n/2 \end{matrix}\right)(y) = \\ &= {}_2F_1\left(\begin{matrix} \rho+\lambda & \rho-\lambda \\ n/2 \end{matrix}\right)\left(\frac{x}{x-1}\right) = (1-x)^{\rho+\lambda} {}_2F_1\left(\begin{matrix} \rho+\lambda & \lambda-\rho + \frac{n}{2} \\ n/2 \end{matrix}\right)(x) \end{aligned}$$

with eigenvalue

$$(4.6) \quad -\frac{\tilde{\Delta}}{4} \phi_\lambda = (\rho^2 - \lambda^2) \phi_\lambda.$$

Note that $\rho = 1/2$ for the unit disk. We are interested in the Weyl calculus acting on H_ν , where $D_{\mathbb{C}}$ is the complexification of D . By [7, 12.18] Z has the complexification $Z_{\mathbb{C}}$, of rank $r_{\mathbb{C}}$ and the half sum of positive roots ρ , given by the following table

Z	$Z_{\mathbb{C}}$	$r_{\mathbb{C}}$	ρ
\mathbb{R}^m	\mathbb{C}^m	1	$\frac{m-1}{4}$
\mathbb{C}^m	$\mathbb{C}^m \times \overline{\mathbb{C}^m}$	2	$\frac{m}{2}$
\mathbb{H}^m	$\mathbb{C}^{2 \times 2m}$	2	$m + \frac{1}{2}$
\mathbb{O}^2	\mathbb{C}_V^{16}	2	$\frac{11}{2}$

Here \mathbb{C}_V^{16} denotes the 16-dimensional exceptional Jordan triple not of tube type. For rank 1 domains we have $X = \mathbb{R}$ and $\Omega = (0, \infty)$. Hence Γ_Ω is the usual Γ -function and $c_\Omega = 1$. Since $n_X = 1$, $n_Y = a-1$, $n_V = a(m-1) = n-a$, $2\rho + 1 = (n+a)/2$, Proposition 3.1 shows

$$\int_D d\mu_0(\zeta) f(\zeta) = 2^{2-a} \frac{\Gamma\left(\nu + 1 - \frac{a}{2}\right)}{\Gamma(\nu - 2\rho)\Gamma\left(\frac{n}{2}\right)} \int_G dg f(g(0))$$

where, for K -invariant functions, we have

$$(4.7) \quad 2^{2-a} \int_G dg f(g(0)) = \int_0^1 dx f(x^{1/2} e) \cdot (1-x)^{-2\rho-1} x^{n/2-1}.$$

Specializing Proposition 3.2 yields the normalizing constant

$$(4.8) \quad c_\nu = 2^{-\nu} {}_2F_1 \left(\begin{matrix} \nu - 2\rho & \nu \\ \nu + 1 - \frac{a}{2} \end{matrix} \right) \left(\frac{1}{2} \right)$$

since $n_Y + n_V/2 = a - 1 + \frac{n-a}{2} = 2\rho$. The main result of this paper, leading to the eigenvalues of the Weyl transform, is the following:

THEOREM 4.1. *Let D be a real bounded symmetric domain of rank 1 and dimension n . Then the Weyl calculus ω_ν satisfies*

$$\begin{aligned} (\omega_\nu^* e_\lambda)(0) &= \frac{\Gamma(\nu - \rho + \lambda) \Gamma(\nu - \rho - \lambda)}{\Gamma(\nu - 2\rho) \Gamma(\nu)} \frac{{}_2F_1 \left(\begin{matrix} \nu - \rho + \lambda & \nu - \rho - \lambda \\ \nu + \frac{n}{2} - 2\rho \end{matrix} \right) \left(\frac{1}{2} \right)}{{}_2F_1 \left(\begin{matrix} \nu - 2\rho & \nu \\ \nu + \frac{n}{2} - 2\rho \end{matrix} \right) \left(\frac{1}{2} \right)} = \\ &= \frac{\Gamma(\nu - \rho + \lambda) \Gamma(\nu - \rho - \lambda)}{\Gamma(\nu - 2\rho) \Gamma(\nu)} \frac{{}_2F_1 \left(\begin{matrix} \frac{n}{2} - \rho + \lambda & \frac{n}{2} - \rho - \lambda \\ \nu + \frac{n}{2} - 2\rho \end{matrix} \right) \left(\frac{1}{2} \right)}{{}_2F_1 \left(\begin{matrix} \frac{n}{2} - 2\rho & \frac{n}{2} \\ \nu + \frac{n}{2} - 2\rho \end{matrix} \right) \left(\frac{1}{2} \right)}. \end{aligned}$$

PROOF. Using (4.7), (4.5) and Lemma 3.2, and applying [8, § 20.2, p. 399, (6)] to the admissible parameters $\alpha = \rho + \lambda$, $\beta = \lambda - \rho + \frac{n}{2}$, $\gamma = \frac{n}{2}$, $\rho = \nu + \lambda - \rho$, $\sigma = \nu$ and $z = -1$, we obtain

$$\begin{aligned} 2^{2-a} \int_G dg \phi_\lambda(g(0)) \mathfrak{b}_\nu(g(0)) &= \int_0^1 dx \phi_\lambda(x^{1/2} e) \mathfrak{b}_\nu(x^{1/2} e) (1-x)^{-2\rho-1} x^{n/2-1} = \\ &= \int_0^1 dx (1-x)^{\rho+\lambda} {}_2F_1 \left(\begin{matrix} \rho + \lambda & \rho - \lambda + \frac{n}{2} \\ n/2 \end{matrix} \right) (x) \left(\frac{1-x}{1+x} \right)^\nu (1-x)^{-2\rho-1} x^{n/2-1} = \\ &= \int_0^1 dx {}_2F_1 \left(\begin{matrix} \rho + \lambda & \lambda - \rho + \frac{n}{2} \\ n/2 \end{matrix} \right) (x) (1-x)^{\nu+\lambda-\rho-1} (1+x)^{-\nu} x^{n/2-1} = \\ &= \frac{\Gamma \left(\frac{n}{2} \right) \Gamma(\nu - \rho + \lambda) \Gamma(\nu - \rho - \lambda)}{\Gamma \left(\nu + \frac{n}{2} - 2\rho \right) \Gamma(\nu)} 2^{-\nu} {}_3F_2 \left(\begin{matrix} \nu - \rho + \lambda & \nu - \rho - \lambda & \nu \\ \nu + \frac{n}{2} - 2\rho & \nu \end{matrix} \right) \left(\frac{1}{2} \right), \end{aligned}$$

with the ${}_3F_2$ -function reducing to ${}_2F_1$. This implies the assertion, since $(\omega_\nu^* e_\lambda)(0)$ is a multiple of this integral normalized at $\lambda = \rho$. \square

REFERENCES

- [1] J. ARAZY - H. UPMEIER, *Invariant symbolic calculi and eigenvalues of invariant operators on symmetric domains*. Conference on *Function Spaces, Interpolation Theory, and related topics in honour of Jaak Peetre on his 65th birthday*. University of Lund, Sweden, August 17-22 2000.
- [2] J. ARAZY - H. UPMEIER, *Covariant symbolic calculi on real symmetric domains*. International Workshop on *Operator Theory and Applications* (Faro, September 12-15 2000). IWOTA-Portugal 2000.
- [3] G. VAN DIJK - M. PEVZNER, *Berezin kernels and tube domains*. J. Funct. Anal., to appear.
- [4] J. FARAUT - A. KORÁNYI, *Analysis on Symmetric Cones*. Clarendon Press, Oxford 1994.
- [5] G. FOLLAND, *Harmonic Analysis in Phase Space*. Princeton Univ. Press, 1989.
- [6] S. HELGASON, *Groups and Geometric Analysis*. Academic Press, New York 1984.
- [7] O. LOOS, *Bounded Symmetric Domains and Jordan Pairs*. Univ. of California, Irvine 1977.
- [8] W. MAGNUS - F. OBERHETTINGER - R.P. SONI, *Formulas and Theorems for the Special Functions of Mathematical Physics*. Springer-Verlag, New York 1966.
- [9] Y. NÉRETIN, *Matrix analogs of Beta-integral and Plancherel formula for Berezin kernel representations*. Preprint.
- [10] A. UNTERBERGER - J. UNTERBERGER, *La série discrète de $SL(2, \mathbb{R})$ et les opérateurs pseudo-différentiels sur une demi-droite*. Ann. Sci. Ec. Norm. Sup., 17, 1984, 83-116.
- [11] A. UNTERBERGER - J. UNTERBERGER, *A quantization of the Cartan domain $BD I(q = 2)$ and operators on the light cone*. J. Funct. Anal., 72, 1987, 279-319.
- [12] A. UNTERBERGER - H. UPMEIER, *The Berezin transform and invariant differential operators*. Comm. Math. Phys., 164, 1994, 563-597.
- [13] H. UPMEIER, *Symmetric Banach Manifolds and Jordan C^* -Algebras*. North Holland 1985.
- [14] H. UPMEIER, *Weyl quantization of symmetric spaces: hyperbolic matrix domains*. J. Funct. Anal., 96, 1991, 297-330.
- [15] G. ZHANG, *Berezin transform on real bounded symmetric domains*. Preprint.

J. Arazy:
 Department of Mathematics
 University of Haifa
 HAIFA 31905 (Israele)
 jarazy@math2.haifa.ac.il

H. Upmeier:
 Fachbereich Mathematik
 Universität Marburg
 35032 MARBURG (Germania)
 upmeier@mathematik.uni-marburg.de